

# An extension theorem with an application to formal tree series

Stephen L. Bloom\*

*Department of Computer Science  
Stevens Institute of Technology  
Hoboken, NJ 07030*

Zoltán Ésik†

*Institute for Informatics  
University of Szeged  
Szeged, Hungary*

## Abstract

Suppose that  $T$  is a grove theory, and  $M$  is a Conway or iteration matrix theory which is a subtheory of  $T$ . We provide a sufficient condition under which the dagger operation on  $M$  can be extended to  $T$  so that  $T$  becomes a Conway or iteration theory. Moreover, we prove a general Kleene type theorem applicable to all grove theories which are Conway theories. We discuss applications to synchronization trees and formal tree series.

## 1 Introduction

In many areas of theoretical and applied computer science, one is interested in solving systems of fixed point equations:

$$\begin{aligned}x_1 &= t_1(x_1, \dots, x_n, y_1, \dots, y_p) \\ &\vdots \\ x_n &= t_n(x_1, \dots, x_n, y_1, \dots, y_p).\end{aligned}\tag{1}$$

where, for each  $i \in [n]$ ,  $t_i$  is a function  $t_i : A^{n+p} \rightarrow A$ , for some structure  $A$ . The structure may be a poset of some kind, and the functions may be order-preserving or continuous. Of course, there are other possible settings in which fixed points are of

---

\*Supported in part by NSF grant 0119916.

†Supported in part by BRICS, Denmark, NSF grant 0119916, and by grant no. T35163 from the National Foundation of Hungary for Scientific Research.

interest: one is when  $t_i$  is a row matrix with  $n + p$  entries in some semiring,  $S$ . One example from language theory is when  $S$  is the idempotent semiring of all subsets of a free monoid  $\Sigma^*$ , with union as sum and complex concatenation as product. Another, is when  $t_i$  is a pair  $(a, v)$ , consisting of a 1 by  $n + p$  row matrix over a semiring  $S$ , and an element  $v$  in an  $S$ -module. Continuing the language example, with the same semiring, the power set of  $\Sigma^*$ , the module might be the collection of all subsets of infinite words on  $\Sigma$ . Another setting is when  $t_i$  is a functor  $\mathcal{D}^{n+p} \rightarrow \mathcal{D}$ , where  $\mathcal{D}$  is some category.

Depending on the context, some questions about such systems are

- Does the system have any solution?
- Is there a canonical (least, initial, etc.) solution?
- Is there a unique solution?
- If there are solutions, what do all solutions look like?
- What are the properties of solutions?

We take as our framework for discussing fixed point equations the categories known as (one-sorted) *Lawvere algebraic theories* [33], categories whose objects are nonnegative integers, and  $n + p$  is the coproduct of  $n, p$ . In the case that the  $t_i$  denote order preserving functions  $A^{n+p} \rightarrow A^n$  on a poset  $A$ , a function  $A^p \rightarrow A^n$  is a morphism  $n \rightarrow p$  (note the reversal of direction to translate from products to coproducts). In the matrix example, an  $n \times p$  matrix over  $S$  becomes a morphism  $n \rightarrow p$ .

Now, in an algebraic theory, we interpret the entire right side of the system of fixed point equations (1) as a morphism  $t : n \rightarrow n + p$ , since each  $t_i$  determines a morphism  $1 \rightarrow n + p$  and the coproduct property produces a corresponding morphism  $t = \langle t_1, \dots, t_n \rangle : n \rightarrow n + p$ . Thus, a *solution* of the system is itself a morphism  $\xi : n \rightarrow p$  which satisfies the “fixed point equation”

$$\xi = t \cdot \langle \xi, \mathbf{1}_p \rangle.$$

If we write a solution  $\xi$  as  $\langle \xi_1, \dots, \xi_n \rangle$ , then, for a theory of functions on powers of a set  $A$ , each  $\xi_i$  is a function  $A^{n+p} \rightarrow A$  and

$$\begin{aligned} \xi_1(y) &= t_1(\xi_1(y), \xi_2(y), \dots, \xi_n(y), y) \\ &\vdots \\ \xi_n(y) &= t_n(\xi_1(y), \xi_2(y), \dots, \xi_n(y), y), \end{aligned}$$

where we write  $y$  for  $y_1, \dots, y_p$ .

Algebraic theories  $T$  in which every system of fixed point equations has a canonical solution are modeled as theories enriched by a function  $\dagger : T(n, n+p) \rightarrow T(n, p)$ , for each  $n, p \geq 0$ , such that, for each morphism  $t : n \rightarrow n+p$ ,

$$t^\dagger = t \cdot \langle t^\dagger, \mathbf{1}_p \rangle.$$

Thus, the operation  $t \mapsto t^\dagger$  produces a canonical solution to the system determined by the morphism  $t$ .

There has been a good deal of effort by both mathematicians and computer scientists devoted to the question of whether fixed points to certain systems exist at all. The names Brouwer [15], Banach [1], Tarski [44], Kleene [26], and Scott [40] come to mind. The *properties* of the fixed point operation in theories which occur naturally in the theory of computation have been studied both in the setting of language theory ([26, 38, 39, 16, 29]) and in setting of domains and continuous functions ([2, 37, 34, 36]). One example of an equational property (other than the fixed point property itself) of the fixed point operation is the “composition identity”:

$$(f \cdot \langle g, 0_n \oplus \mathbf{1}_p \rangle)^\dagger = f \cdot \langle (g \cdot \langle f, 0_m \oplus \mathbf{1}_p \rangle)^\dagger, \mathbf{1}_p \rangle,$$

all  $f : n \rightarrow m+p$ ,  $g : m \rightarrow n+p$ . As a result of a long series of papers, there seems to be agreement that there is essentially one equational theory which captures all of the equations valid in every computationally interesting theory. These equations may be described in any of several equivalent ways:

- all equations valid in theories of complete partially ordered sets and continuous functions;
- all equations valid in theories of continuous functors on  $\omega$ -complete categories;
- all equations valid in theories of labelled trees;
- all models of a certain set of axioms.

The models for these equations are called *iteration theories*. The reason for the qualification “essentially one” above is that certain iteration theories, such as a theory of matrices over the semiring of binary relations on some set, are special only in that they have a unique morphism  $1 \rightarrow 0$ . Thus, in addition to the axioms mentioned, they also satisfy  $x = y$ , all  $x, y : 1 \rightarrow 0$ .

The best set of axioms known for iteration theories is due to Ésik, [25]. His axioms are divided into two parts: the “Conway identities”, which contain only two axiom

schemes, the parameter and double dagger identities, in addition to the composition identity above, and an identity for each finite (simple) group. Each of the axioms is readily seen to hold in any particular setting. However, the completeness of these axioms is not at all obvious.

The parameter identity is

$$(f \cdot (\mathbf{1}_n \oplus g))^\dagger = f^\dagger \cdot g,$$

for all  $f : n \rightarrow n + p$ ,  $g : p \rightarrow q$ . The double dagger identity is

$$f^{\dagger\dagger} = (f \cdot (\langle \mathbf{1}_n, \mathbf{1}_n \rangle \oplus \mathbf{1}_p))^\dagger,$$

all  $f : n \rightarrow n + n + p$ . (A brief account of theories and the axiomatization iteration theories is given in the appendix.)

There is a class of iteration theories in which the “nontrivial” systems of fixed point equations have unique solutions: these theories were introduced by Elgot in [22] under the term “(ideal) iterative theories”. In such a theory, a morphism  $1 \rightarrow n$  either is one of the coproduct injections  $i_n : 1 \rightarrow n$ ,  $i \in [n]$ , or is *ideal*; if  $f : 1 \rightarrow n$  is ideal, so is  $f \cdot g$ , for any  $g : n \rightarrow p$ . A morphism  $f : n \rightarrow p$  is ideal if each component  $i_n \cdot f$  is. For each ideal morphism  $f : n \rightarrow n + p$  in an iterative theory, there is a unique  $\xi = f^\dagger$  which is a solution to the “Elgot fixed point equation”

$$\xi = f \cdot \langle \xi, \mathbf{1}_p \rangle. \quad (2)$$

One class of examples of ideal iterative theories are the “contraction theories”. If  $A$  is a complete metric space, the least theory containing all proper contractions  $A^n \rightarrow A$  is an iterative theory.

One might legitimately consider fixed point equations for ideal morphisms as equations in which the variables to solve for occur only in “guarded positions”. Thus, iterative theories might be described roughly as theories in which systems of guarded equations have unique fixed points. In iterative theories, the nonideal morphisms are somehow trivial, and can be dealt with as in [5, 6]. Models in which “guarded statements” of the required form have unique fixed points have been featured in many places, for example in the metric semantics of de Bakker and his collaborators, see, e.g., [18, 19, 20], in the work of Milner, e.g., [35], and in the classical paper axiomatizing the regular sets, by Salomaa, [38].

What about morphisms  $n \rightarrow n + p$  in iterative theories which are not ideal? For example, consider the case  $n = 1$ ,  $p = 0$  and  $f = \mathbf{1}_1$ , the identity morphism. The corresponding fixed point equation is the trivial equation in the variable  $\xi : 1 \rightarrow 0$ ,

$$\xi = \mathbf{1}_1 \cdot \xi,$$

which has all morphisms  $1 \rightarrow 0$  as solutions. In [5, 6] it is shown that for each choice of a solution to this one fixed point equation, there is a unique extension of the fixed point operation to all morphisms such that the resulting theory is an iteration theory.

The theme of extending the fixed point operation from subtheories in which the solution is unique to the entire theory - while preserving its properties - was continued in several settings: for matrix theories in [8], and for matricial theories in [9]. In a matrix theory  $T$ , the hom-set  $T(n, p)$  is a commutative monoid and composition distributes over the monoid operation on both sides. A matrix theory is isomorphic to one in which a morphism  $n \rightarrow p$  is an  $n \times p$  matrix over a fixed semiring, the semiring of morphisms  $1 \rightarrow 1$ . As an application of the matrix extension theorem in [8], it was shown that if the theory of matrices over the semiring  $S$  is a Conway or an iteration theory, then so is the theory of matrices over the semiring of all formal power series over  $S$ . In Section 3 we review that result.

In the current paper, we continue playing the same tune, but in a different setting. We consider what we have called “grove theories” in [7], motivated by an analogous use of the term in [3]. A grove theory is a relaxation of the notion of a matrix theory: composition distributes over the monoid operation only on one side.

The outline of this paper is the following. After reviewing the extension theorem for matrix theories in Section 3, we define grove theories in Section 4, and state and prove the new extension theorem in Sections 5 and 6.

We then consider two important classes of examples. One example involves a different kind of tree: the synchronization trees. In Section 7, we use the extension theorem to show that the theory of synchronization trees is an iteration theory. It then follows that synchronization trees modulo bisimulation equivalence and other behavioral equivalences form iteration theories. This result is known, but our original argument (see [7]) made use of 2-categories and continuity arguments.

In Section 8, we discuss a new example in more detail, the theory of **formal tree series** [4, 32] over a semiring. We first formulate this theory and then use the grove extension theorem to show that if the semiring  $S$  is a Conway or iteration semiring, then for each signature  $\Sigma$ , so is the theory of formal  $\Sigma$ -trees over  $S$ .

In Section 9, we state and prove a general Kleene theorem, applicable to all Conway grove theories. In the following Section 10, we apply this Kleene theorem to the theory of formal tree series, after introducing the appropriate notion of “rational expression” for grove theories.

The Appendix (Sections 11, 12, 13) contains a quick review of theories and iteration

theories.

A preliminary version of this paper appeared in [11].

## 2 Preliminaries

We denote the set  $\{1, 2, \dots, n\}$  by  $[n]$ . The composite of morphisms  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is written in diagrammatic order:

$$f \cdot g = A \xrightarrow{f} B \xrightarrow{g} C.$$

We give a review of Lawvere algebraic theories and various tree theories in the Appendix. In particular, the axioms for Conway and iteration theories are presented there.

## 3 The Matrix Extension Theorem

A **matrix theory**  $T$  is a pointed theory, i.e., a theory with a distinguished morphism  $0_{1,0} : 1 \rightarrow 0$ , such that each hom-set  $T(n, p)$  is a commutative monoid, with neutral

element  $0_{n,p} = \overbrace{\langle 0_{1,0}, \dots, 0_{1,0} \rangle}^n \cdot 0_p$  such that

$$\begin{aligned} (f + g) \cdot h &= f \cdot h + g \cdot h \\ k \cdot (f + g) &= k \cdot f + k \cdot g, \\ 0_{r,n} \cdot f &= 0_{r,p} \\ f \cdot 0_{p,q} &= 0_{n,q} \end{aligned}$$

for all  $f, g : n \rightarrow p$ ,  $h : p \rightarrow q$  and  $k : r \rightarrow n$  in  $T$ . A **morphism**  $T \rightarrow T'$  of matrix theories is a theory morphism which preserves the monoid structure on each hom-set (and thus preserves the point).

In [23] it is shown that if  $T$  is a matrix theory, then  $T$  is isomorphic to a theory of matrices  $\mathbf{Mat}_S$  over the semiring  $S = T(1, 1)$ . A morphism  $n \rightarrow p$  in  $\mathbf{Mat}_S$  is an  $n \times p$  matrix over  $S$ . Composition of morphisms is matrix multiplication, and the additive structure is defined in the expected way.

In matrix theories which are preiteration theories, one may define a “star” operation on the square matrices by

$$a^* := [a, \mathbf{1}_n]^\dagger,$$

where  $a$  is  $n \rightarrow n$ . Using this operation, the Conway identities take the form of some familiar identities for regular expressions, which is the reason for the terminology. The double dagger identity becomes the “star sum” identity:

$$(a + b)^* = (a^*b)^*a^*, \quad (3)$$

where  $a, b : n \rightarrow n$ . The composition identity becomes the “product star” identity:

$$(ab)^* = \mathbf{1}_n + a(ba)^*b, \quad (4)$$

where  $a : n \rightarrow m$  and  $b : m \rightarrow n$ . The parameter identity holds automatically. See [16] and [8]. Also, in any Conway matrix theory, the dagger operation is related to the star operation by:

$$[a, b]^\dagger = a^*b,$$

when  $a : n \rightarrow n$  and  $b : n \rightarrow p$ . A **Conway**, respectively **iteration matrix theory** is a matrix theory which is a Conway theory, respectively, an iteration theory. A **morphism** of Conway (or iteration) matrix theories is both a matrix theory morphism and a Conway theory morphism. If  $S$  is a semiring such that  $\mathbf{Mat}_S$  is a Conway or iteration theory, then we call  $S$  a **Conway**, or an **iteration semiring**. The collection of all Conway (or iteration) semirings forms a variety of algebras whose signature is that of semirings enriched with a unary star operation. See [7].

In [8], the following “extension theorem” was proved. Suppose that  $S$  is a subsemiring of the semiring  $R$ . Suppose further that there is an ideal  $I$  of  $R$  such that  $R$  is the direct sum of  $S$  and  $I$ , in the sense that any element  $x$  of  $R$  can be written uniquely as  $x = s + a$ , where  $s \in S$  and  $a \in I$ .

**Theorem 3.1 (Matrix Extension Theorem)** *If  $\mathbf{Mat}_S$  is a Conway (resp. iteration) theory, and if for each  $a \in I, b \in R$  there is a unique  $\xi \in R$  such that  $\xi = a\xi + b$  in  $R$ , then, there is a unique extension of the dagger operation on  $\mathbf{Mat}_S$  to  $\mathbf{Mat}_R$  so that  $\mathbf{Mat}_R$  becomes a Conway (resp. iteration) theory.*

It follows from this result that if the theory of matrices over the semiring  $S$  is a Conway or iteration theory, then so is the theory of matrices over the semiring of formal power series over  $S$  with variables in the set  $X$ .

Indeed, a formal power series  $r$  over the semiring  $S$  with variables in  $X$  is a function  $r : X^* \rightarrow S$ . In this case, the semiring  $S$  is isomorphic to the collection of functions  $r : X^* \rightarrow S$  such that  $r(u) = 0$  for all nonempty words  $u \in X^*$ , and the ideal  $I$  is the collection of all functions  $a : X^* \rightarrow S$  with  $a(\lambda) = 0$ , where  $\lambda$  is the word of length zero.

## 4 Grove theories

A **grove theory**  $T$  is a pointed theory such that each hom-set  $T(n, p) = (T(n, p), +, 0_{n,p})$  is a commutative monoid with neutral element  $0_{n,p}$  such that

$$\begin{aligned} (f + g) \cdot h &= f \cdot h + g \cdot h \\ 0_{r,n} \cdot f &= 0_{r,p} \\ i_n \cdot (f + g) &= i_n \cdot f + i_n \cdot g, \quad i \in [n] \\ i_n \cdot 0_{n,p} &= 0_{1,p}, \quad i \in [n], \end{aligned}$$

for all  $f, g : n \rightarrow p, h : p \rightarrow q$  in  $T$ . (The last two equations say that sums of vector morphisms are defined componentwise, as are the neutral elements.) If  $T, T'$  are grove theories, a **morphism**  $\varphi : T \rightarrow T'$  is a theory morphism that is a monoid morphism  $T(n, p) \rightarrow T'(n, p)$ , for each  $n, p \geq 0$ .  $T$  is a **sub grove theory** of  $T'$  if  $T$  is a subtheory of  $T'$ , and the monoid  $T(n, p)$  is a submonoid of  $T'(n, p)$ , for each  $n, p \geq 0$ .

Each matrix theory is a grove theory, but not conversely.

In any grove theory, for a nonnegative integer  $n$ , define the morphism  $\mathbf{n} : 1 \rightarrow 1$  as

$$\mathbf{n} := \overbrace{\mathbf{1}_1 + \dots + \mathbf{1}_1}^n.$$

The following fact is easy to prove by induction.

**Proposition 4.1** *In any grove theory  $T$ , and any nonnegative integer  $n$ ,*

$$\begin{aligned} \mathbf{n} \cdot (f + g) &= \mathbf{n} \cdot f + \mathbf{n} \cdot g \\ \mathbf{n} \cdot 0_{1,p} &= 0_{1,p}, \end{aligned}$$

*all  $f, g : 1 \rightarrow p$  in  $T$ .*

Suppose that  $T$  is a grove theory and  $T_0$  is a subtheory of  $T$ . We say that  $T_0$  is additively closed if  $T_0(n, p)$  is a submonoid of  $T(n, p)$ , for each  $n, p \geq 0$ .

**Corollary 4.2** *The matrix theory  $\mathbf{Mat}_{\mathbb{N}}$  of matrices over the semiring of nonnegative integers is initial in the class of grove theories and matrix theories.*

A **Conway** or **iteration grove theory** is a grove theory which is also a Conway or iteration theory. Thus, any Conway or iteration matrix theory is a Conway or

iteration grove theory. A **morphism** of Conway or iteration grove theories is a grove theory morphism which is also a Conway or iteration theory morphism.

We let  $\mathbf{N}_\infty$  denote the semiring consisting of the nonnegative integers with an “infinite” element  $\infty$  adjoined; the addition and multiplication are defined in the expected way. This semiring is a Conway and iteration semiring when the star operation is defined by

$$x^* := \begin{cases} 1 & x = 0 \\ \infty & \text{otherwise.} \end{cases}$$

The corresponding matrix theory is a Conway and iteration theory. In  $\mathbf{Mat}_{\mathbf{N}_\infty}$ , the equation  $1^* = 2^*$  holds; in the dagger version,

$$(1_2 + 2_2)^\dagger = (\mathbf{2} \cdot 1_2 + 2_2)^\dagger. \quad (5)$$

**Proposition 4.3** *The matrix theory  $\mathbf{Mat}_{\mathbf{N}_\infty}$  is initial in the class of Conway and iteration grove theories satisfying (5) It is also initial in the class of Conway and iteration matrix theories satisfying this equation.*

We omit the proof. For the last claim, see also Corollary 5.5 in [7], Chapter 9.

We now turn to the statement of the main result.

## 5 The grove extension theorem

The hypotheses take a while to state, but they are similar to those needed for the matrix extension theorem. First we need a definition.

**Definition 5.1** *Suppose that  $T$  is a grove theory and  $M$  is a sub grove theory of  $T$  which is a matrix theory. A collection  $T_0$  of morphisms of  $T$  is an  **$M$ -ideal** if  $T_0$*

- *contains all zero morphisms  $0_{n,p}$ ;*
- *is closed under addition, i.e., if  $f, g : n \rightarrow p \in T_0$ , then  $f + g \in T_0$ ;*
- *is closed under right composition, i.e., if  $f : n \rightarrow p$  is in  $T_0$  and  $g : p \rightarrow q$  is any morphism, then  $f \cdot g : n \rightarrow q$  is in  $T_0$ ;*
- *is closed under left composition with any morphism in  $M$ .*

### 5.1 The hypotheses

Assume that  $T$  is a grove theory and  $M$  is a sub grove theory of  $T$  that is a matrix theory. Further, we assume that

H1.  $T_0$  is an  $M$ -ideal of  $T$ .

H2. Every  $f : n \rightarrow p$  in  $T$  can be written uniquely as

$$f = a + f_0$$

for some  $a \in M$ ,  $f_0 \in T_0$ . We say:  $T$  is the *direct sum* of  $M$  and  $T_0$ .

H3. If  $a \in M$  and  $f, g \in T$  have appropriate sources and targets, then

$$a \cdot (f + g) = a \cdot f + a \cdot g.$$

H4. For every  $f_0 : n \rightarrow n + p$  in  $T_0$  and  $b : n \rightarrow p$  in  $M$ , there is a unique solution  $\xi : n \rightarrow p$  to the equation

$$\xi = ((0_n \oplus b) + f_0) \cdot \langle \xi, \mathbf{1}_p \rangle,$$

and the solution is written

$$((0_n \oplus b) + f_0)^\nabla.$$

H5.  $M$  is a Conway theory, so that there is a  $*$ -operation on the square matrices satisfying

$$\begin{aligned} (ab)^* &= \mathbf{1}_n + a(ba)^*b, & a : n \rightarrow m, & b : m \rightarrow n \\ (a+b)^* &= (a^*b)^*a^*, & a, b : n \rightarrow n. \end{aligned}$$

In particular,  $S = M(1, 1)$  is a Conway semiring.

**Theorem 5.2** [*The grove extension theorem*] *Under the above hypotheses, there is a unique way to define a dagger operation*

$$\dagger : T(n, n + p) \longrightarrow T(n, p),$$

*on each hom-set  $T(n, n + p)$  extending that on  $M$  such that  $T$  becomes a Conway theory. Further, if  $M$  satisfies the weak functorial implication, so does  $T$ ; if  $M$  is an iteration theory, so is  $T$ .*

The proof of this theorem is in Section 6. The “weak functorial implication” is defined in the Appendix.

We can revise the assumptions needed for the grove extension theorem to apply only to “scalar morphisms”, i.e., those with source 1.

**Proposition 5.3** *Under the assumptions that  $T$  is a grove theory,  $S \subseteq T(1,1)$  is a Conway semiring (which determines the matrix subtheory  $M = \mathbf{Mat}_S$  of  $T$ ), and  $I_0$  is a collection of scalar morphisms of  $T$ , the hypotheses H1–H5 are equivalent to the hypotheses H1’–H4’ below.*

H1’a. If  $s \in S$ ,  $f : 1 \rightarrow p \in I_0$ ,  $g : p \rightarrow q \in T$ , then  $s \cdot f \cdot g \in I_0$ .

H1’b. If  $f, g : 1 \rightarrow p \in I_0$ , then  $f + g \in I_0$ .

H1’c.  $0_{1,p} \in I_0$ , all  $p \geq 0$ .

H2’. For any  $g : 1 \rightarrow p$  in  $T$  there is a unique  $a : 1 \rightarrow p \in M$  and  $f : 1 \rightarrow p \in I_0$  such that  $g = f + a$ .

H3’. For any  $f, g : 1 \rightarrow p \in T$  and any  $s \in S$ ,  $s \cdot (f + g) = s \cdot f + s \cdot g$ .

H4’. For any  $b : 1 \rightarrow p \in M$  and any  $f : 1 \rightarrow 1 + p$  in  $I_0$ , there is a unique  $\xi : 1 \rightarrow p \in T$  such that

$$\xi = ((0_1 \oplus b) + f) \cdot \langle \xi, \mathbf{1}_p \rangle.$$

Then the hypotheses H1–H5 above hold, and conversely.

*Proof sketch.* It is clear that H1–H5 imply H1’–H4’, by letting  $I_0$  be the collection of all morphisms  $T_0(1, p)$ ,  $p \geq 0$ . Now, assume that H1’–H4’ hold. We define  $T_0(n, p)$  as the collection of all morphisms  $f = \langle f_1, \dots, f_n \rangle$  such that  $f_i \in I_0$ , for each  $i \in [n]$ . Then, the only point which may not be clear is that for  $b : n \rightarrow p$  in  $\mathbf{Mat}_S$ , and  $f : n \rightarrow n + p$  in  $T_0$ , there is a unique solution to the Elgot fixed point equation (2) for  $(0_n \oplus b) + f$ . But this fact follows from the inductive argument in [12], which shows that “scalar iterative theories” are also iterative theories. This shows that the hypotheses of the grove extension theorem hold.  $\square$

## 6 Proof of the main theorem

In this section, we prove the grove extension theorem, Theorem 5.2.

**Uniqueness.** First, we show that there is at most one such extension.

Suppose that  $T$  has such a dagger operation making it a Conway theory. If  $g = ((0_n \oplus b) + f_0)$ , for  $f_0 : n \rightarrow n + p$  in  $T_0$  and  $b : n \rightarrow p$  in  $M$ , then, by the fixed point identity,

$$g^\dagger = ((0_n \oplus b) + f_0) \cdot \langle g^\dagger, \mathbf{1}_p \rangle.$$

Thus

$$((0_n \oplus b) + f_0)^\dagger = ((0_n \oplus b) + f_0)^\nabla.$$

So any fixed point operation  $\dagger$  must extend  $\nabla$ .

We prove a useful proposition.

**Proposition 6.1** *Assume that  $T$  is a Conway grove theory with a subtheory  $M$  which is a Conway matrix theory. Then, for any  $f_0 : n \rightarrow n + p$  in  $T$  and  $[a, b] : n \rightarrow n + p$  in  $M$ ,*

$$(f_0 + [a, b])^\dagger = (a^* \cdot ((0_n \oplus b) + f_0))^\dagger.$$

**Lemma 6.2** *With the assumptions of Proposition 6.1 on  $T$  and  $M$ , if  $a : n \rightarrow n$  in  $M$  and  $g : n \rightarrow n + p$  in  $T$ , then*

$$((a \oplus 0_{n+p}) + (0_n \oplus g))^\dagger = a^* g : n \longrightarrow n + p$$

*Proof.* Since  $\dagger$  extends the dagger operation on  $M$ ,

$$\begin{aligned} ((a \oplus 0_{n+p}) + (0_n \oplus \mathbf{1}_n \oplus 0_p))^\dagger &= [a, \mathbf{1}_n, 0_{n,p}]^\dagger \\ &= [a^*, 0_{n,p}] \\ &= a^* \oplus 0_p. \end{aligned}$$

Then, since

$$((a \oplus 0_{n+p}) + (0_n \oplus g)) = ((a \oplus 0_{n+p}) + (0_n \oplus \mathbf{1}_n \oplus 0_p)) \cdot (\mathbf{1}_n \oplus \langle g, 0_n \oplus \mathbf{1}_p \rangle),$$

we have, using the parameter identity,

$$\begin{aligned} ((a \oplus 0_{n+p}) + (0_n \oplus g))^\dagger &= ((a \oplus 0_{n+p}) + (0_n \oplus \mathbf{1}_n \oplus 0_p))^\dagger \cdot \langle g, 0_n \oplus \mathbf{1}_p \rangle \\ &= (a^* \oplus 0_p) \cdot \langle g, 0_n \oplus \mathbf{1}_p \rangle \\ &= a^* \cdot g. \quad \square \end{aligned}$$

We now complete the proof of Proposition 6.1. Suppose that  $f = ([a, b] + f_0) : n \rightarrow n + p$  in  $T$ ,  $f_0 \in T$ , and  $[a, b] : n \rightarrow n + p$  in  $M$ . Then

$$f = (a \oplus 0_p) + ((0_n \oplus b) + f_0).$$

If  $g = ((0_n \oplus b) + f_0)$ , then,

$$\begin{aligned} f^\dagger &= ((a \oplus 0_p) + g)^\dagger \\ &= (((a \oplus 0_{n+p}) + (0_n \oplus g)) \cdot (\langle \mathbf{1}_n, \mathbf{1}_n \rangle \oplus \mathbf{1}_p))^\dagger \\ &= ((a \oplus 0_{n+p}) + (0_n \oplus g))^{\dagger\dagger}, \end{aligned}$$

using the double dagger identity,

$$= (a^* \cdot g)^\dagger,$$

by Lemma 6. This fact proves Proposition 6.1.

Now if  $f_0 \in T_0$ , then, by Proposition 6.1,

$$(a^* \cdot g)^\dagger = (a^* \cdot g)^\nabla.$$

Indeed, since  $T_0$  is an  $M$ -ideal,  $a^* \cdot f_0 \in T_0$ . Also,

$$\begin{aligned} a^* \cdot g &= a^* \cdot ((0_n \oplus b) + f_0) \\ &= (0_n \oplus a^*b) + (a^* \cdot f_0), \end{aligned}$$

by H3. Thus, since  $\dagger$  extends  $\nabla$ , we have  $(a^* \cdot g)^\dagger = (a^* \cdot g)^\nabla$ . This shows that  $(f_0 + [a, b])^\dagger = (a^*g)^\nabla$ , so that  $\dagger$  is determined by  $*$  and  $\nabla$ .  $\square$

### Existence.

We use the formula derived above to define  $\dagger$ . So, given  $f = ([a, b] + f_0) : n \rightarrow n + p$ , with  $[a, b] \in M$  and  $f_0 \in T_0$ , we **define**

$$f^\dagger := (a^* \cdot g)^\nabla,$$

where

$$g := (0_n \oplus b) + f_0 : n \longrightarrow n + p.$$

Since  $f$  can be written as  $[a, b] + f_0$  in exactly one way,  $f \mapsto f^\dagger$  is well defined.

**Lemma 6.3** *The dagger operation extends the dagger operation on  $M$ .*

*Proof.* Choose  $f_0 = 0_{n,n+p} \in T_0$ . Then, by definition, if  $f = [a, b] + f_0 = [a, b] : n \rightarrow n + p$ , then

$$g = (0_n \oplus b) + f_0 = 0_n \oplus b,$$

so that

$$\begin{aligned} a^* \cdot g &= 0_n \oplus a^*b, \quad \text{and} \\ f^\dagger &= (0_n \oplus a^*b)^\nabla \\ &= a^*b, \end{aligned}$$

which is the dagger operation in  $M$ . □

**Lemma 6.4** *The dagger operation extends  $\nabla$ .*

Suppose that  $f = (0_p \oplus b) + f_0$ , so that  $a = 0_{n,n}$ . Then, since  $0_{n,n}^* = \mathbf{1}_n$ ,

$$\begin{aligned} f^\dagger &= (0_{n,n}^* \cdot f)^\nabla \\ &= f^\nabla. \quad \square \end{aligned}$$

From now on, we drop the use of  $\nabla$ , so that

$$((a \oplus 0_p) + f)^\dagger = (a^* \cdot f)^\dagger,$$

whenever  $a : n \rightarrow n \in M$  and  $f = (0_n \oplus b) + f_0 : n \rightarrow n + p$ , with  $b : n \rightarrow p$  in  $M$  and  $f_0 \in T_0$ .

To save space, we write  $f \in \hat{T}_0$  if  $f = (0_n \oplus b) + f_0 : n \rightarrow n + p$ , as above.

**Lemma 6.5** *For  $a : n \rightarrow n$  in  $M$  and  $f : n \rightarrow n + p \in \hat{T}_0$ ,*

$$(a^* \cdot f)^\dagger = a^* \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger.$$

*Thus,*

$$((a \oplus 0_p) + f)^\dagger = a^* \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger.$$

*Proof.* By hypotheses H1 and H3,  $a^* \cdot f \in \hat{T}_0$ . Thus, it is enough to show that  $a^* \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger$  is a solution of the fixed point equation for  $a^* \cdot f$ .

$$\begin{aligned} a^* \cdot f \cdot \langle a^* \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger, \mathbf{1}_p \rangle &= a^* \cdot f \cdot (a^* \oplus \mathbf{1}_p) \cdot \langle (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger, \mathbf{1}_p \rangle \\ &= a^* \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger, \end{aligned}$$

since the fixed point identity holds for morphisms in  $\hat{T}_0$ .

In order to show that  $T$  is a Conway theory when equipped with this dagger operation, we will show that it satisfies the parameter, composition and double dagger identities (see [7], Chapter 6.2). For use in the argument, we show first that it satisfies the fixed point identity.

**Proposition 6.6** *The operation  $\dagger$  satisfies the fixed point identity.*

*Proof.* Suppose that  $a : n \rightarrow n$  in  $M$  and  $f : n \rightarrow n + p$  in  $\hat{T}_0$ . We want to show

$$((a \oplus 0_p) + f)^\dagger = ((a \oplus 0_p) + f) \cdot \langle ((a \oplus 0_p) + f)^\dagger, \mathbf{1}_p \rangle,$$

in other words, that

$$([a, 0] + f) \cdot \langle (a^* \cdot f)^\dagger, \mathbf{1}_p \rangle = (a^* \cdot f)^\dagger,$$

where we write just 0 for  $0_{np}$ . But,

$$([a, 0] + f) \cdot \langle (a^* \cdot f)^\dagger, \mathbf{1}_p \rangle = a(a^* \cdot f)^\dagger + f \cdot \langle (a^* \cdot f)^\dagger, \mathbf{1}_p \rangle,$$

since composition on the right distributes over sums,

$$= a(a^* \cdot f)^\dagger + f \cdot \langle a^* \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger, \mathbf{1}_p \rangle$$

by Lemma 6.5,

$$\begin{aligned} &= a(a^* \cdot f)^\dagger + f \cdot (a^* \oplus \mathbf{1}_p) \cdot \langle (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger, \mathbf{1}_p \rangle \\ &= a(a^* \cdot f)^\dagger + (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger \\ &= aa^* \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger + (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger \\ &= (aa^* + \mathbf{1}_n) \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger \\ &= a^* \cdot (f \cdot (a^* \oplus \mathbf{1}_p))^\dagger \\ &= (a^* \cdot f)^\dagger, \end{aligned}$$

again, by Lemma 6.5. Note that the morphism  $f \cdot (a^* \oplus \mathbf{1}_p)$  belongs to  $\hat{T}_0$ . We used the fact that by assumptions H5 and H4, the fixed point identity holds for morphisms in  $M$  and  $\hat{T}_0$ .

**Proposition 6.7** *The composition identity holds.*

*Proof.* For notational simplicity, we prove this in the parameter free case. We will show that

$$((a + f) \cdot (b + g))^\dagger = (a + f) \cdot ((b + g) \cdot (a + f))^\dagger, \quad (6)$$

where  $a : n \rightarrow m$ ,  $b : m \rightarrow n$  in  $M$  and  $f : n \rightarrow m$ ,  $g : m \rightarrow n$  in  $\hat{T}_0$ . The left hand side of (6) is

$$((ab)^*(f \cdot (b + g) + ag))^\dagger$$

and since the morphism  $(ab)^*(f \cdot (b + g) + ag)$  is in  $\hat{T}_0$ , it is enough to show that the right hand side of (6) is a solution of the iteration equation for this morphism.

Let  $h$  abbreviate the morphism  $((b + g)(a + f))^\dagger$ , so that

$$h = ((ba)^*(g \cdot (a + f) + bf))^\dagger.$$

Then,

$$\begin{aligned} (ab)^*(f \cdot (b + g) + ag)(a + f) \cdot h &= (ab)^*f \cdot (b + g)(a + f) \cdot h \\ &\quad + (ab)^*ag(a + f) \cdot h \\ &= (ab)^*f \cdot h + (ab)^*ag(a + f) \cdot h \\ &= a(ba)^*bf \cdot h + f \cdot h + a(ba)^*g(a + f) \cdot h \\ &= a(ba)^*(bf + g(a + f))((ba)^*(bf + g(a + f)))^\dagger \\ &\quad + f \cdot h \\ &= a((ba)^*(bf + g(a + f)))^\dagger + f \cdot h \\ &= a \cdot h + f \cdot h \\ &= (a + f)h. \quad \square \end{aligned}$$

**Proposition 6.8** *The parameter identity holds.*

First, we need the following fact.

**Lemma 6.9** *If  $f : n \rightarrow n + p$  is in  $\hat{T}_0$ , so is  $f \cdot (\mathbf{1}_n \oplus g)$ , for any  $g : p \rightarrow q$ .*

*Proof.* Indeed, if  $f = (0_n \oplus b) + f_0$ , as always, then  $f_0 \cdot (\mathbf{1}_n \oplus g)$  is in  $T_0$ , since  $T_0$  is closed under right composition with any morphism. Also,

$$\begin{aligned} f \cdot (\mathbf{1}_n \oplus g) &= ((0_n \oplus b) + f_0) \cdot (\mathbf{1}_n \oplus g) \\ &= (0_n \oplus (bg)) + (f_0 \cdot (\mathbf{1}_n \oplus g)). \end{aligned}$$

Now, writing  $g = (a + g_0)$ , where  $a : p \rightarrow q$  in  $M$  and  $g_0 : p \rightarrow q$  in  $T_0$ , we have

$$\begin{aligned} bg &= b \cdot (a + g_0) \\ &= b \cdot a + b \cdot g_0, \end{aligned}$$

by H3. Now  $b \cdot a \in M$  and  $b \cdot g_0 \in T_0$ . Thus, we have

$$\begin{aligned} f \cdot (\mathbf{1}_n \oplus g) &= (0_n \oplus b') + f'_0, \quad \text{where} \\ b' &= b \cdot a \\ f'_0 &= b \cdot g_0 + f_0 \cdot (\mathbf{1}_n \oplus g), \end{aligned}$$

completing the proof.  $\square$

Now suppose that  $a : n \rightarrow n$  in  $M$ ,  $f : n \rightarrow n + p$  in  $\hat{T}_0$  and  $g : p \rightarrow q$  in  $T$ . We want to show that

$$((([a, 0] + f) \cdot (\mathbf{1}_n \oplus g)))^\dagger = (([a, 0] + f)^\dagger \cdot g.$$

But

$$\begin{aligned} ((([a, 0] + f) \cdot (\mathbf{1}_n \oplus g)))^\dagger &= ([a, 0] + f \cdot (\mathbf{1}_n \oplus g))^\dagger \\ &= (a^* \cdot f \cdot (\mathbf{1}_n \oplus g))^\dagger. \end{aligned}$$

Now, since the parameter identity always holds for unique fixed points, ([7], Chapter 5) and since the morphism  $a^* \cdot f \cdot (\mathbf{1}_n \oplus g)$  is in  $\hat{T}_0$ ,

$$\begin{aligned} (a^* \cdot f \cdot (\mathbf{1}_n \oplus g))^\dagger &= (a^* \cdot f)^\dagger \cdot g \\ &= (([a, 0] + f)^\dagger \cdot g, \end{aligned}$$

as was to be shown.  $\square$

We now turn to the double dagger identity.

**Proposition 6.10** *The double dagger identity holds.*

*Proof.* We consider only the parameter free case. Let  $g = [a, b] + f : n \rightarrow n + n$ , where  $[a, b] \in M$  and  $f \in T_0$ . Let  $\rho = \langle \mathbf{1}_n, \mathbf{1}_n \rangle$ . We want to show that

$$(g \cdot \rho)^\dagger = g^{\dagger\dagger}. \quad (7)$$

But,

$$\begin{aligned} (g \cdot \rho)^\dagger &= (([a, b] + f) \cdot \rho)^\dagger \\ &= (a + b + (f \cdot \rho))^\dagger \\ &= ((a + b)^* \cdot (f \cdot \rho))^\dagger. \end{aligned}$$

Suppose that we can prove

$$((a+b)^* \cdot f \cdot \rho) \cdot g^{\dagger\dagger} = g^{\dagger\dagger}, \quad (8)$$

then, since  $(a+b)^* \cdot (f \cdot \rho) \in T_0$ , we have

$$((a+b)^* \cdot f \cdot \rho)^{\dagger} = g^{\dagger\dagger},$$

showing that (7) holds.

So we now prove (8).

$$\begin{aligned} g^{\dagger} &= ([a, b] + f)^{\dagger} \\ &= ([a, 0] + ([0, b] + f))^{\dagger} \\ &= (a^*([0, b] + f))^{\dagger} \\ &= ([0, a^*b] + a^* \cdot f)^{\dagger}. \end{aligned}$$

But, by the fixed point identity proved above,

$$\begin{aligned} ([0, a^*b] + a^* \cdot f)^{\dagger} &= ([0, a^*b] + a^* \cdot f) \cdot \langle ([0, a^*b] + a^* \cdot f)^{\dagger}, \mathbf{1}_n \rangle \\ &= ([0, a^*b] + a^* \cdot f) \cdot \langle g^{\dagger}, \mathbf{1}_n \rangle \\ &= a^*b + a^* \cdot f \cdot \langle g^{\dagger}, \mathbf{1}_n \rangle. \end{aligned}$$

Thus,

$$\begin{aligned} g^{\dagger\dagger} &= ((a^*b)^* a^* \cdot f \cdot \langle g^{\dagger}, \mathbf{1}_n \rangle)^{\dagger} \\ &= ((a+b)^* \cdot f \cdot \langle g^{\dagger}, \mathbf{1}_n \rangle)^{\dagger}. \end{aligned} \quad (9)$$

We now complete the proof of (8).

$$\begin{aligned} (a+b)^* \cdot f \cdot \rho \cdot g^{\dagger\dagger} &= (a+b)^* \cdot f \cdot \langle g^{\dagger\dagger}, g^{\dagger\dagger} \rangle \\ &= (a+b)^* \cdot f \cdot \langle g^{\dagger}, \mathbf{1}_n \rangle \cdot g^{\dagger\dagger}, \end{aligned}$$

by the fixed point identity for  $g^{\dagger}$ ,

$$\begin{aligned} &= (a+b)^* \cdot f \cdot \langle g^{\dagger}, \mathbf{1}_n \rangle \cdot ((a+b)^* \cdot f \cdot \langle g^{\dagger}, \mathbf{1}_n \rangle)^{\dagger} \\ &= ((a+b)^* \cdot f \cdot \langle g^{\dagger}, \mathbf{1}_n \rangle)^{\dagger} \\ &= g^{\dagger\dagger}, \end{aligned}$$

by two applications of (9). □

This completes the proof that with the extension,  $T$  is a Conway theory.

**Proposition 6.11** *Suppose that the weak functorial implication holds in  $M$ . Then it also holds in  $T$ .*

*Proof.* Suppose that the diagram below commutes, where  $\rho : n \rightarrow 1$  is the unique base morphism,  $a : n \rightarrow n$ ,  $b : 1 \rightarrow 1$  in  $M$ , and  $f, g$  are in  $\hat{T}_0$ .

$$\begin{array}{ccc} n & \xrightarrow{(a \oplus 0_p) + f} & n + p \\ \rho \downarrow & & \downarrow \rho \oplus \mathbf{1}_p \\ 1 & \xrightarrow{(b \oplus 0_p) + g} & 1 + p \end{array}$$

Since  $T$  is the direct sum of  $M$  and  $T_0$ , it follows that

$$\begin{aligned} a \cdot \rho &= \rho \cdot b \\ f \cdot (\rho \oplus \mathbf{1}_p) &= \rho \cdot g. \end{aligned}$$

Since the weak functorial implication holds in  $M$ ,

$$a^* \cdot \rho = \rho \cdot b^*.$$

Since the weak functorial implication holds for unique fixed points,

$$f^\dagger = \rho \cdot g^\dagger.$$

In fact,

$$\begin{aligned} a^* \cdot f \cdot (\rho \oplus \mathbf{1}_p) &= a^* \cdot \rho \cdot g \\ &= \rho \cdot b^* \cdot g, \end{aligned}$$

so that

$$(a^* \cdot f)^\dagger = \rho \cdot (b^* \cdot g)^\dagger,$$

completing the proof. □

Last, we show that if  $M$  is an iteration theory, so is  $T$ .

**Proposition 6.12** *If  $M$  satisfies the group equation associated with the finite group  $G$ , then so does  $T$ .*

*Proof.* Let  $h : 1 \rightarrow n + p$  be any morphism and suppose that  $G$  is a group of order  $n$  (whose underlying set is  $[n]$ ). Let  $h_G$  denote the morphism  $n \rightarrow n + p$ :

$$\langle h \cdot (\rho_1^G \oplus \mathbf{1}_p), \dots, h \cdot (\rho_n^G \oplus \mathbf{1}_p) \rangle$$

where the morphisms  $\rho_i^G : n \rightarrow n$  are the base morphisms determined by the group operation  $i, j \mapsto i \bullet j$  of  $G$ :

$$\rho_i^G(j) := i \bullet j.$$

We use the same notation for morphisms in  $M$ : If  $a = [a_1, \dots, a_n] : 1 \rightarrow n$  in  $M$ ,

$$a_G = \begin{bmatrix} a_{\rho_1^G(1)} & \cdots & a_{\rho_1^G(n)} \\ \vdots & \ddots & \vdots \\ a_{\rho_n^G(1)} & \cdots & a_{\rho_n^G(n)} \end{bmatrix}$$

It is known from [29] that

$$(a_G)^* = b_G,$$

for some  $b = [b_1, \dots, b_n]$  in  $M$ .

So assume that  $a : 1 \rightarrow n$  in  $M$  and  $f : 1 \rightarrow n + p$  in  $\hat{T}_0$ . Let  $\bar{a}$  denote the sum of the entries in  $a$ , so that  $\bar{a} = a \cdot \rho$ , where  $\rho : n \rightarrow 1$  is base. Let  $\bar{f} = f \cdot (\rho \oplus \mathbf{1}_p)$ .

We want to show that

$$((a_G \oplus 0_p) + f_G)^\dagger = \rho \cdot ((\bar{a} \oplus 0_p) + \bar{f})^\dagger. \quad (10)$$

The left hand side is

$$((a_G)^* f_G)^\dagger = (b_G \cdot f_G)^\dagger.$$

But  $b_G \cdot f_G \in \hat{T}_0$ , and

$$\begin{aligned} b_G \cdot f_G \cdot \rho &= b_G \cdot \rho \cdot \bar{f} \\ &= \rho \cdot \bar{b} \cdot \bar{f}, \end{aligned}$$

where  $\bar{b}$  is the sum of the entries of  $b_G$ . Since the weak functorial implication holds for unique fixed points ([7], Chapter 5), we have

$$(b_G \cdot f_G)^\dagger = \rho \cdot (\bar{b} \cdot \bar{f})^\dagger,$$

i.e.,

$$((a_G)^* f_G)^\dagger = \rho \cdot (\bar{b} \cdot \bar{f})^\dagger.$$

But since the group identity holds in  $M$ ,  $\bar{b}$  is just  $\bar{a}^*$ . Thus,

$$\begin{aligned} ((a_G)^* f_G)^\dagger &= \rho(\bar{a}^* \bar{f})^\dagger \\ &= \rho \cdot ((\bar{a} \oplus 0_p) + \bar{f})^\dagger, \end{aligned}$$

proving (10). □

The proof of the grove extension theorem is complete.

We now discuss two applications of the theorem.

## 7 The theory of synchronization trees

Let  $A$  be a nonempty set. The theory  $ST(A)$  of **synchronization trees over  $A$**  has been studied in [7]. We recall that a morphism  $1 \rightarrow p$  in this theory is a tree  $t = (V, v_0, E, \ell)$ , where

1.  $V$  is a finite or countable set;
2.  $(V, v_0, E)$  is a rooted tree - i.e.,  $E \subseteq V \times V$ ,  $v_0 \in V$  and, for each  $v \in V$  there is a unique path  $v_0, v_1, \dots, v_k = v$  from  $v_0$  to  $v$ .
3.  $\ell : E \rightarrow A \cup [p]$  is a “labeling function”.
4. if  $\ell(u, v) \in [p]$ , then  $v$  is a leaf, and  $(u, v)$  is an “exit edge”.

A morphism  $s : p \rightarrow q$  is a  $p$ -tuple of trees  $1 \rightarrow q$ . If  $t : 1 \rightarrow p$  and  $s = (s_1, \dots, s_p)$  are synchronization trees, the **composite**  $t \cdot s$  is the tree obtained from  $t$  by replacing any exit edge labeled  $i \in [p]$  by a copy of the tree  $s_i$ ; if  $t, s : 1 \rightarrow p$  are synchronization trees,  $t + s : 1 \rightarrow p$  is the tree obtained from the disjoint union of the two sets of vertices and edges by identifying the roots of  $s, t$  and otherwise making no further identifications. Composition and sum is extended to  $n$ -tuples of trees pointwise. (See [7] for all details.)

The tree  $0_{1,p} : 1 \rightarrow p$  is the unique tree with only one vertex, the root, and no edges. The morphism  $0_{n,p}$  is the  $n$ -tuple of the trees  $0_{1,p}$ .

In [7], it was shown that  $ST(A)$  is a grove theory.

**Proposition 7.1** *Each collection of trees  $n \rightarrow p$  in  $ST(A)$  is a complete metric space, where the distance between two trees  $n \rightarrow p$  is the maximum of the distances between their components  $i_n \cdot t, i_n \cdot t'$ . For trees  $t, t' : 1 \rightarrow p$ , define the distance  $d(t, t')$  to be 0 if  $t = t'$ , or  $1/2^n$ , where  $n$  is the minimum depth for which the trees  $t|n$  and  $t'|n$  are distinct. The tree  $t|n$  is  $t$  “cut off” at level  $n$ ; prune all vertices and edges at distance more than  $n$  from the root.*

Let  $Ex$  be the subtheory of  $ST(A)$  determined by the trees  $1 \rightarrow p$  all of whose edges are exit edges; thus, for each  $i$  in  $[p]$ , there are either finitely many exit edges labeled  $i$  or countably many such edges. Recall that  $\mathbf{N}_\infty$  denotes the semiring of the nonnegative integers with a point at infinity adjoined. For any synchronization tree  $t : 1 \rightarrow p$  in  $Ex$ , let  $m_i$  denote the number of edges in the tree labeled  $i$  whose source is the root, for  $i \in [p]$ . We may thus identify such a tree  $t$  with the  $1 \times p$  matrix  $m(t) = [m_1, \dots, m_p]$  over  $\mathbf{N}_\infty$ . This identification respects both the theory sum and composition operations, as is easy to show. Thus,  $Ex$  is isomorphic to the theory  $\mathbf{Mat}_{\mathbf{N}_\infty}$  of all matrices over the semiring  $\mathbf{N}_\infty$ .

Now let  $T_0$  be the collection of all morphisms  $f = \langle f_1, \dots, f_n \rangle : n \rightarrow p$  in  $ST(A)$  such that each component  $f_i$  is a tree with the following property. No edge whose source is the root is labeled by an integer in  $[p]$ . These trees have been called “guarded” in [7].

It is straightforward to check that the hypotheses listed in Proposition 5.3 hold. In particular, if  $f : n \rightarrow n + p$  is in  $T_0$  and  $b : n \rightarrow p$  is in  $Ex$ , then there is a unique tree which is a solution of the iteration equation for  $(0_n \oplus b) + f : n \rightarrow n + p$ , namely the equation

$$\xi = ((0_n \oplus b) + f) \cdot \langle \xi, \mathbf{1}_p \rangle.$$

Indeed, the tree  $(0_n \oplus b) + f$  induces a proper contraction map  $C$  on the set of trees  $t : n \rightarrow p$  in  $ST(A)$ , where

$$C(t) := ((0_n \oplus b) + f) \cdot \langle t, \mathbf{1}_p \rangle.$$

The distance between  $C(t)$  and  $C(t')$  is at most  $1/2$  the distance between  $t$  and  $t'$ . Thus, by the Banach fixed point theorem [1],  $C$  has a unique fixed point.

Thus, as a corollary to the grove extension theorem, we obtain

**Theorem 7.2** *There is a unique way to define  $\dagger$  on  $ST(A)$  extending the corresponding operation on  $Ex$ , so that  $ST(A)$  is a Conway theory. And since  $Ex$  is in fact an iteration theory, so is  $ST(A)$ . Moreover, the weak functorial implication holds in  $ST(A)$ .*

The fact that  $ST(A)$  is an iteration theory was proved in a totally different way in [7], using 2-categorical notions, initiality and continuity arguments.

## 8 Theories of formal tree series

The second class of examples, apparently a new one, are the grove theories of formal tree series. Recall that  $T_\Sigma(X_n)$  denotes the set of finite  $\Sigma$ -trees whose leaves are labeled by letters in  $X_n \cup \Sigma_0$  and whose interior nodes with  $k$  successors are labeled by letters in  $\Sigma_k$ . The *height*,  $\text{ht}(t)$ , of a tree  $t$  in  $T_\Sigma(X_n)$  is defined as usual.

Assume  $S$  is a *commutative* semiring.

**Definition 8.1** A *formal  $\Sigma$ -tree series*  $s : 1 \rightarrow p$  *over*  $S$  is a function

$$s : T_\Sigma(X_p) \longrightarrow S.$$

We let  $S\langle\langle \Sigma^p \rangle\rangle$  denote the set of all formal  $\Sigma$ -tree series  $1 \rightarrow p$  over  $S$ . We write the value of  $s \in S\langle\langle \Sigma^p \rangle\rangle$  on the tree  $t$  as  $(s, t)$ . The **support** of  $s$  is  $\{t \in T_\Sigma(X_p) : (s, t) \neq 0\}$ .

The following fact is well-known.

**Proposition 8.2** The set  $S\langle\langle \Sigma^p \rangle\rangle$  is a complete metric space, where for  $s, s' \in S\langle\langle \Sigma^p \rangle\rangle$ , the distance function is defined as follows.

$$d(s, s') := \begin{cases} 0 & s = s' \\ 1/2^n & n = \min\{\text{ht}(t) : (s, t) \neq (s', t)\}. \end{cases}$$

Each set  $S\langle\langle \Sigma^p \rangle\rangle$  may be enriched with a commutative addition operation, defined by:

$$(s_1 + s_2, t) := (s_1, t) + (s_2, t), \quad t \in T_\Sigma(X_p).$$

**Proposition 8.3**  $(S\langle\langle \Sigma^p \rangle\rangle, +, \mathbf{0}_p)$  is a commutative monoid, where  $\mathbf{0}_p$  is the constant function with value  $0 \in S$ .  $\square$

**Definition 8.4** For any tree  $t \in T_\Sigma(X_p)$ , let  $\eta(t) \in S\langle\langle \Sigma^p \rangle\rangle$  be the series defined by:

$$(\eta(t), u) := \begin{cases} 1 & u = t \\ 0 & \text{otherwise.} \end{cases}$$

We will need the following construction below.

**Definition 8.5** Suppose that  $t \in T_\Sigma(X_n)$  has exactly  $k$  leaves labeled by variables in  $X_n$ . (There may be other leaves labeled by letters in  $\Sigma_0$ .) Say these leaves are labeled, from left to right, by  $x_{i_1}, \dots, x_{i_k}$ . Let  $t'$  in  $T_\Sigma(X_k)$  be the tree obtained from  $t$  by relabeling these leaves  $x_1, \dots, x_k$ , respectively, and let  $f$  denote the function  $f : [k] \rightarrow [n]$  which maps  $j \in [k]$  to  $i_j \in [n]$ , (and the corresponding base morphism  $k \rightarrow n$ ). Then

$$t = 1 \xrightarrow{t'} k \xrightarrow{f} n.$$

When  $k = 0$ ,  $t'$  is the same tree as  $t$ , and  $f = 0_n$ .

**Definition 8.6** For any tree  $u \in T_\Sigma(X_p)$ , let  $D_n(u)$  denote the set of all trees  $t \in T_\Sigma(X_n)$  such that

$$u = t' \cdot \langle u_1, \dots, u_k \rangle$$

for some trees  $u_1, \dots, u_k \in T_\Sigma(X_p)$ , where  $t'$  is constructed from  $t$  as in Definition 8.5.

We note that  $D_n(u)$  is a finite set, which is nonempty when  $n > 0$ . In particular, for  $i \in [p]$ ,

$$D_n(x_i) = \{x_1, \dots, x_n\}, \quad (11)$$

since if  $x_i = t' \cdot \langle u_1, \dots, u_k \rangle$ , we have  $k = 1$ ,  $t' = x_1$  and  $u_1 = x_i$ . Thus, if  $t = x_j$ ,  $f(1) = j$ . Further, for a fixed  $u \in T_\Sigma(X_p)$  and  $t \in D_n(u)$ , there is a unique set of trees  $u_1, \dots, u_k$  such that  $u = t' \cdot \langle u_1, \dots, u_k \rangle$ .

We define next the **theory**  $S\langle\langle \Sigma \rangle\rangle$  of formal  $\Sigma$ -tree series over  $S$ .

**Definition 8.7** A morphism  $1 \rightarrow p$  in  $S\langle\langle \Sigma \rangle\rangle$  is a function in  $S\langle\langle \Sigma^p \rangle\rangle$ ; a morphism  $n \rightarrow p$  is an  $n$ -tuple  $\langle s_1, \dots, s_n \rangle$  of morphisms  $1 \rightarrow p$ . For  $s \in S\langle\langle \Sigma^n \rangle\rangle$ ,  $s_i \in S\langle\langle \Sigma^p \rangle\rangle$ ,  $i \in [n]$ , and  $u \in T_\Sigma(X_p)$ , we **define**

$$(s \cdot \langle s_1, \dots, s_n \rangle, u) := \sum_{t \in D_n(u)} (s, t) \cdot (s_{f(1)}, u_1) \cdots (s_{f(k)}, u_k), \quad (12)$$

where  $t = t' \cdot f$ , as in Definition 8.5, and  $D_n(u)$  is defined in Definition 8.6, and  $u, t', u_1, \dots, u_k$  are related as in that Definition. The composite of  $\langle s_1, \dots, s_n \rangle$  with  $r : n \rightarrow q$  is defined by:

$$\langle s_1, \dots, s_n \rangle \cdot r := \langle s_1 \cdot r, \dots, s_n \cdot r \rangle.$$

For  $i \in [n]$ , the coproduct injection  $i_n : 1 \rightarrow n$  is the series  $\eta(x_i)$  in  $S\langle\langle \Sigma^n \rangle\rangle$ .

When the semiring is the Boolean semiring, this composition is known as the “OI” substitution, [24].

In order to make  $S\langle\langle \Sigma \rangle\rangle$  into a grove theory, we extend the addition operation to all hom-sets pointwise: for morphisms  $s, r : n \rightarrow p$  in  $S\langle\langle \Sigma \rangle\rangle$ , with  $s = \langle s_1, \dots, s_n \rangle$  and  $r = \langle r_1, \dots, r_n \rangle$ , we **define**

$$\begin{aligned} s + r &:= \langle s_1 + r_1, \dots, s_n + r_n \rangle \\ \mathbf{0}_{n,p} &:= \langle \mathbf{0}_p, \dots, \mathbf{0}_p \rangle. \end{aligned}$$

We omit the proof that  $S\langle\langle \Sigma \rangle\rangle$  is a theory, and note only

**Proposition 8.8**  $S\langle\langle \Sigma \rangle\rangle$  is a grove theory.

Indeed, since the sum of two  $n$ -tuples of morphisms  $1 \rightarrow p$  is defined pointwise, we need only check that  $S\langle\langle \Sigma^p \rangle\rangle$  is a commutative monoid, and composition on the right distributes over sum. Both facts are easy to verify.  $\square$

We will use the next fact several times.

**Lemma 8.9** Suppose  $s : 1 \rightarrow n$ ,  $s_i : 1 \rightarrow p$ , in  $S\langle\langle \Sigma \rangle\rangle$ , for each  $i \in [n]$ . Then, for each variable  $x_j \in X_p$ ,

$$(s \cdot \langle s_1, \dots, s_n \rangle, x_j) = \sum_{i \in [n]} (s, x_i) \cdot (s_i, x_j).$$

*Proof.* Indeed, by (12),

$$(s \cdot \langle s_1, \dots, s_n \rangle, x_j) = \sum_{t \in D_n(x_j)} (s, t) \cdot (s_{f(1)}, u_1) \cdots (s_{f(k)}, u_k),$$

where  $f, k$  are related to  $t$  by Definition 8.5. But then  $t \in X_n$ , by (8.6), so that  $k = 1$ ,  $t' = x_1$ , and  $u_1 = x_j$ . If  $t = x_i$ ,  $f(1) = i$  and

$$(s \cdot \langle s_1, \dots, s_n \rangle, x_j) = \sum_{i \in [n]} (s, x_i) \cdot (s_i, x_j). \quad \square$$

For any  $g = \langle s_1, \dots, s_n \rangle : n \rightarrow p$  in  $S\langle\langle \Sigma \rangle\rangle$ , let  $\mu(g)$  denote the  $n \times p$  matrix in  $\mathbf{Mat}_S$ , such that

$$\mu(g)_{i,j} := (s_i, x_j).$$

**Proposition 8.10** For  $f : n \rightarrow p$  and  $g : p \rightarrow q$  in  $S\langle\langle \Sigma \rangle\rangle$ ,

$$\mu(f \cdot g) = \mu(f) \cdot \mu(g).$$

If  $f, g : n \rightarrow p$  in  $S\langle\langle \Sigma \rangle\rangle$ ,

$$\mu(f + g) = \mu(f) + \mu(g).$$

*Proof.* It is enough to prove both statements for the case  $n = 1$ . The result for composition follows from Lemma 8.9, and for addition by definition of  $f + g$ .  $\square$

Now let  $M(n, p)$  consist of all morphisms  $\langle s_1, \dots, s_n \rangle : n \rightarrow p$  in  $S\langle\langle \Sigma \rangle\rangle$  such that, for each  $i \in [n]$ , the support of  $s_i$  is a subset of  $X_p$ .

**Proposition 8.11**  $M$  is a sub grove theory of  $S\langle\langle \Sigma \rangle\rangle$ , i.e.,  $\mathbf{0}_{n,p} \in M(n, p)$ ; if  $g_1, g_2 \in M(n, p)$ , then  $g_1 + g_2 \in M(n, p)$ , and if  $h \in M(p, q)$ ,  $g_1 \cdot h \in M(n, q)$ , for each  $n, p, q \geq 0$ . Also,  $\eta(x_i) \in M(1, p)$ , for each  $i \in [p]$ .

*Proof.* It is enough to prove these facts when  $n = 1$ . Now if  $t \notin X_p$ ,  $(g_1 + g_2, t) = (g_1, t) + (g_2, t) = 0$ , so that  $M(1, p)$  is closed under addition. To show that  $M$  is closed under composition, write  $g_1 = s$ ,  $h = \langle s_1, \dots, s_p \rangle$ . Then, for any  $u \in T_\Sigma(X_q)$ ,

$$(s \cdot h, u) = \sum_{t \in D_n(u)} (s, t) \cdot (s_{f(1)}, u_1) \cdots (s_{f(k)}, u_k),$$

as usual. But unless  $t \in X_n$ , the value  $(s, t)$  is 0. Thus, for  $t = x_i \in X_n$ ,  $t' = x_1$ ,  $k = 1$ ,  $f(1) = i$  and  $u = u_1$ , so that  $(s \cdot h, u) = \sum_{i \in [n]} (s, x_i) \cdot (s_i, u)$ . But unless  $u \in X_q$ , we have  $(s_i, u) = 0$ . Thus, the support of  $s \cdot h$  is a subset of  $X_q$ .  $\square$

**Corollary 8.12**  $M$  is isomorphic to the matrix theory  $\mathbf{Mat}_S$ .

*Proof.* This follows from Proposition 8.10 and from the fact that the restriction of  $\mu$  to  $M$  is a bijection  $M \rightarrow \mathbf{Mat}_S$ .  $\square$

**Proposition 8.13** If  $a : n \rightarrow p$  in  $M$  and  $s, s' : p \rightarrow q$  in  $S\langle\langle \Sigma \rangle\rangle$ , then

$$a \cdot (s + s') = a \cdot s + a \cdot s'.$$

*Proof.* We assume  $n = 1$ . Since the support of  $a$  is a subset of  $X_p$ , for any  $u \in T_\Sigma(X_q)$ , and any  $r = \langle r_1, \dots, r_p \rangle : p \rightarrow q$ , by (12),

$$(a \cdot r, u) = \sum_{i \in [p]} (a, x_i) \cdot (r_i, u).$$

The result follows immediately.  $\square$

Let  $T_0(1, p)$  denote the set of morphisms  $f_0 : 1 \rightarrow p$  in  $S\langle\langle \Sigma \rangle\rangle$  such that

$$i \in [p] \implies (s, x_i) = 0.$$

Let  $T_0(n, p)$  be those morphisms  $n \rightarrow p$  such that each component is in  $T_0(1, p)$ .

We note the following obvious fact.

**Proposition 8.14** *Any  $f : n \rightarrow p$  in  $S\langle\langle \Sigma \rangle\rangle$  can be written uniquely as  $f = a + g$ , where  $a : n \rightarrow p$  in  $M$  and  $g : n \rightarrow p$  in  $T_0$ .*  $\square$

**Proposition 8.15** *Suppose that  $f, g \in T_0(n, p)$ . Then  $f + g \in T_0(n, p)$ . If  $a \in M(k, n)$ , then  $a \cdot f \in T_0(k, p)$ . If  $h : p \rightarrow q$  in  $S\langle\langle \Sigma \rangle\rangle$ , then  $f \cdot h \in T_0(n, q)$ . The constant function in  $S\langle\langle \Sigma^p \rangle\rangle$  with value 0 belongs to  $T_0(1, p)$ .*

*Proof.* We prove closure of  $T_0$  under left composition with morphisms in  $M$ . It is enough to assume  $k = 1$ , so let  $s : 1 \rightarrow n$  in  $M$  and  $g = \langle s_1, \dots, s_n \rangle$  in  $T_0(n, p)$ . By Lemma 8.9, for each  $x_j \in X_q$ ,

$$(s \cdot g, x_j) = \sum_{i \in [n]} (s, x_i) \cdot (s_i, x_j).$$

But, for each  $i \in [n]$ ,  $s_i \in T_0(1, q)$ , so that  $(s_i, x_j) = 0$ . Hence  $(s \cdot g, x_j) = 0$ .

The fact that  $T_0$  is closed under right composition with any morphism is proved in the same way, using Lemma 8.9 again.  $\square$

**Proposition 8.16** *If  $g \in T_0(1, 1 + p)$  and  $b : 1 \rightarrow p$  in  $M$ , then there is a unique  $\xi : 1 \rightarrow p \in S\langle\langle \Sigma \rangle\rangle$  such that*

$$\xi = ((0_1 \oplus b) + g) \cdot \langle \xi, \mathbf{1}_p \rangle.$$

*Proof.* Recall that  $S\langle\langle \Sigma^p \rangle\rangle$  is a complete metric space. We claim that the function  $C : S\langle\langle \Sigma^p \rangle\rangle \rightarrow S\langle\langle \Sigma^p \rangle\rangle$  defined by

$$C(s) := ((0_1 \oplus b) + g) \cdot \langle s, \mathbf{1}_p \rangle$$

is a proper contraction on the complete metric space  $S\langle\langle \Sigma^p \rangle\rangle$ . It then follows from the Banach fixed point theorem that  $C$  has a unique fixed point, which is the statement of the proposition.

Let  $r \in S\langle\langle \Sigma^{1+p} \rangle\rangle$  be defined as

$$r = (0_1 \oplus b) + g.$$

Note that  $\mu(0_1 \oplus b)$  is the matrix

$$[0, (b, x_1), \dots, (b, x_p)].$$

We compute the value of  $r$  on each  $x_j \in X_{1+p}$ .

$$(r, x_j) = \begin{cases} (g, x_1) = 0 & j = 1 \\ (b, x_{j-1}) + (g, x_j) = (b, x_{j-1}) & j > 1 \end{cases}$$

since  $g \in T_0(1, 1+p)$ .

Now  $C(s) = r \cdot \langle s, s_2, \dots, s_{1+p} \rangle$ , where, for  $j \in [p]$ ,  $s_{1+j} = \eta(x_j)$ . Thus, by Lemma 8.9, for  $j \in [p]$ ,

$$\begin{aligned} (C(s), x_j) &= (r, x_1) \cdot (s, x_j) + \sum_{i \in [p]} (b, x_i) \cdot (\eta(x_i), x_j) \\ &= \sum_{i \in [p]} (b, x_i) \cdot (\eta(x_i), x_j) \\ &= (b, x_j). \end{aligned}$$

since  $(r, x_1) = 0$ . For an atomic tree  $u = \sigma_0 \cdot 0_p$ , with  $\sigma_0 \in \Sigma_0$ ,  $(C(s), u) = (g, \sigma_0 \cdot 0_{1+p})$ . Thus, for any  $s_1, s_2$  in  $S\langle\langle \Sigma^p \rangle\rangle$ ,  $C(s_1)$  and  $C(s_2)$  agree on all trees in  $T_\Sigma(X_p)$  of depth 0. For a tree  $u \in T_\Sigma(X_p)$  of depth at least 1,

$$(C(s), u) = \sum_{t \in D_{1+p}(u)} (r, t) \cdot (s_{f(1)}, u_1) \cdots (s_{f(k)}, u_k),$$

where  $u = t' \cdot \langle u_1, \dots, u_k \rangle$ ,  $t = t' \cdot f$ , as always. Let  $A$  be the collection of all trees in  $D_{1+p}(u)$  that have at least one leaf labeled  $x_1$ , and let  $B = D_{1+p}(u) - A$ . Then

let

$$\begin{aligned} C_A(s, u) &= \sum_{t \in A} C_t(s, u) \\ C_B(s, u) &= \sum_{t \in B} C_t(s, u), \end{aligned}$$

where, for  $t \in D_{1+p}(u)$ ,

$$C_t(s, u) := (r, t) \cdot (s_{f(1)}, u_1) \cdots (s_{f(k)}, u_k).$$

Then  $(C(s), u) = C_A(s, u) + C_B(s, u)$ . Note that for *any*  $s, s'$  in  $S\langle\langle \Sigma^p \rangle\rangle$  and any  $t \in B$ ,

$$C_t(s, u) = C_t(s', u),$$

since for trees  $t \in B$ ,  $f(i) > 1$ , for all  $i \in [k]$ , so that the functions  $s_{f(i)}$  are in the set  $\{\eta(x_1), \dots, \eta(x_p)\}$ . Let  $\beta$  be this common value  $C_B(s, u)$ . So,  $(C(s), u) = \beta + C_A(s, u)$ . If  $C(s, u) \neq C(s', u)$ , there must be at least one tree  $t$  in  $A$  with  $C_t(s, u) \neq C_t(s', u)$ . Assume that  $t$  is one such tree. We know  $\text{ht}(t) > 0$ . Assume further, for ease of notation, that  $f^{-1}(1) = \{1, 2, \dots, j\}$ , for some  $j$ ,  $1 \leq j \leq k$ . Then

$$\begin{aligned} C_t(s, u) &= (r, t) \cdot (s, u_1) \cdots (s, u_j) (s_{f(j+1)}, u_{j+1}) \cdots (s_{f(k)}, u_k) \\ C_t(s', u) &= (r, t) \cdot (s', u_1) \cdots (s', u_j) (s_{f(j+1)}, u_{j+1}) \cdots (s_{f(k)}, u_k) \end{aligned}$$

Hence,  $(s, u_i) \neq (s', u_i)$ , for at least one  $i \in [j]$ . Since

$$u = t' \cdot \langle u_1, \dots, u_k \rangle$$

and  $\text{ht}(t') = \text{ht}(t) > 0$ ,

$$d(C(s), C(s')) \leq \frac{1}{2}d(s, s'). \quad \square$$

Since we have verified the hypotheses of the Proposition 5.3, by the grove extension theorem, we get the following result.

**Corollary 8.17** *If  $S$  is a Conway or iteration semiring, there is a unique extension of the fixed point operation on  $M$  to all of  $S\langle\langle \Sigma \rangle\rangle$  so that  $S\langle\langle \Sigma \rangle\rangle$  is a Conway or iteration theory.*

The *proof* of the grove extension theorem gives some more information as to the value of  $s^\dagger$ , for  $s : 1 \rightarrow 1 + p$  in  $S\langle\langle \Sigma \rangle\rangle$ .

Indeed, suppose we write  $s = [a, b] + g$ , where  $[a, b] : 1 \rightarrow 1 + p$  in  $\mathbf{Mat}_S$  and  $g \in T_0$ . Then we know, from this proof, that  $s^\dagger = (a^* \cdot (0_1 \oplus b + g))^\dagger$ . But

$$a^* \cdot ((0_1 \oplus b) + g) = (0_1 \oplus (a^*b)) + (a^* \cdot g),$$

and  $a^* \cdot g$  is in  $T_0$ . Thus, by the Banach fixed point theorem,  $s^\dagger$  is the metric limit  $\lim_n s_n$ , where

$$\begin{aligned} s_0 &:= \mathbf{0}_p \\ s_{n+1} &:= ((0_1 \oplus (a^*b)) + (a^*g)) \cdot \langle s_n, \mathbf{1}_p \rangle \\ &= a^*b + (a^*g) \cdot \langle s_n, \mathbf{1}_p \rangle \end{aligned}$$

We could have chosen any element in  $S\langle\langle \Sigma^p \rangle\rangle$  as  $s_0$ . In the case that  $S$  is a continuous, naturally ordered semiring,  $\mathbf{0}_p$  is the least in  $S\langle\langle \Sigma^p \rangle\rangle$ , so that, with  $s_0 = \mathbf{0}_p$ ,  $s^\dagger = \sup_n s_n$ , as well.  $\square$

## 9 A general Kleene theorem

Assume that  $T$  is a Conway grove theory and  $M$  is a sub Conway grove theory of  $T$  which is a matrix theory. For a given collection  $\Sigma$  of scalar morphisms in  $T$  we will describe the smallest sub Conway grove theory of  $T$  containing all the morphisms in  $M \cup \Sigma$ .

In more detail, we are assuming that

1.  $T$  is a Conway grove theory and  $M$  is a sub Conway theory of  $T$  that is a matrix theory. Thus,  $M$  has a  $*$ -operation on the  $p \times p$  matrices, for each  $p \geq 1$ , satisfying the star sum and product identities (3) and (4) in Section 3 above. The hom-set  $M(1, 1)$  is a Conway semiring, which we denote by  $S$ .
2. If  $a \in M$  and  $f, g \in T$  have appropriate sources and targets, then

$$a \cdot (f + g) = a \cdot f + a \cdot g.$$

Let  $\Sigma$  be a collection of scalar morphisms in  $T$ .

**Definition 9.1** A morphism  $f : 1 \rightarrow n + p$  is **primitive** of weight  $n$  if  $f$  is a finite sum of morphisms of the sort

$$\begin{aligned} s \cdot \sigma \cdot \rho & \quad (\text{type 1}) \\ 0_n \oplus (s \cdot \tau) & \quad (\text{type 2}), \end{aligned}$$

where  $s \in S$ ,  $\sigma : 1 \rightarrow k \in \Sigma$ ,  $\rho : k \rightarrow n + p$  and  $\tau : 1 \rightarrow p$  are base. A morphism  $f = \langle f_1, \dots, f_q \rangle : q \rightarrow n + p$  is **primitive** of weight  $n$  if each component is primitive of weight  $n$ .

Note that any  $1 \times p$  matrix in  $M$  is primitive of type 2, (when  $n = 0$ ), so, in effect, all morphisms in  $M$  are primitive (of type 2).

We list some easy consequences of these definitions.

**Proposition 9.2** 1. If  $f : r \rightarrow n + p$  is primitive of weight  $n$ , so is  $\gamma \cdot f : m \rightarrow n + p$ , for any base morphism  $\gamma : m \rightarrow r$ .

2. If  $f : 1 \rightarrow n + p$  is primitive of weight  $n$  of type 1, then  $f \cdot \gamma : 1 \rightarrow n + q$  is primitive of weight  $n$  of type 1, for any base  $\gamma : n + p \rightarrow n + q$ .

3. If  $f : 1 \rightarrow n + p$  is primitive of weight  $n$  of type 2, then  $f \cdot (\mathbf{1}_n \oplus 0_m \oplus \gamma) : 1 \rightarrow n + m + q$  is primitive of weight  $n + m$  of type 2, for any base  $\gamma : p \rightarrow q$ .

4. If  $g : m \rightarrow n + p$  is primitive of weight  $n$ , and  $b : m \rightarrow m$  is in  $M$ , then  $b \cdot g$  is primitive of weight  $n$ . Indeed, if  $m = 1$ ,  $b = [s]$ , say, and  $g = g_1 + \dots + g_r$ , then  $b \cdot g = s \cdot g_1 + \dots + s \cdot g_r$ , since left composition with  $M$ -morphisms distributes over sums. And if  $g_i$  is primitive of any type, so is  $s \cdot g_i$ .

5. If  $f : m \rightarrow n + p$  is primitive of weight  $n$ , then  $0_q \oplus f$  is primitive of weight  $q + n$ .

6. If  $0_q \oplus f : m \rightarrow q + n + p$  is primitive of weight  $q + n$ , then  $f$  is primitive of weight  $n$ .

**Definition 9.3** A **normal description**  $k \rightarrow p$  of weight  $n$  (over  $(\Sigma, S)$ ) is a pair

$$D = (\alpha, f),$$

where  $f : n \rightarrow n + p$  is primitive of weight  $n$ , and  $\alpha : k \rightarrow n$  is base. The **behavior** of the normal description  $D : k \rightarrow p$  is the morphism

$$|D| := \alpha \cdot f^\dagger.$$

Below, we will often write a primitive morphism  $f$  of weight  $n$  as

$$f = \bar{f} + (0_n \oplus a),$$

where  $\bar{f}$  is of type 1 and  $a : n \rightarrow p \in M$ .

A normal description  $D = (\alpha, f) : k \rightarrow p$  of weight  $n$  might be viewed as an “automaton” with  $n$  internal states, and  $p$  exit states; the values of  $\alpha$  pick out the  $k$  start or final states among the internal states, and the behavior of  $D$  is the morphism in  $T$  determined by this automaton.

The rest of this section contains a proof of the following “Kleene type” theorem.

**Theorem 9.4** *The smallest sub Conway grove theory of  $T$  containing  $M$  and  $\Sigma$  consists of the behaviors of normal descriptions.*

*Proof.* Let  $\mathcal{D}$  denote the collection of all morphisms of the form  $|D|$ , for a normal description  $D$ .

First, it is clear that any sub Conway grove theory of  $T$  that contains  $S \cup \Sigma$  will contain each morphism in  $\mathcal{D}$ . Thus, we need show only that this collection of morphisms is itself a sub Conway grove theory of  $T$ .

**$\mathcal{D}$  is closed under composition.**

Suppose that  $D = (\alpha, f) : k \rightarrow p$  has weight  $n$  and  $E = (\beta, g) : p \rightarrow q$  has weight  $m$ . Write

$$\begin{aligned} f &= \bar{f} + (0_n \oplus a) \\ g &= \bar{g} + (0_m \oplus b), \end{aligned}$$

where  $a : n \rightarrow p$  and  $b : m \rightarrow q$  in  $M$ . Then, define the normal description  $D \cdot E$  as

$$D \cdot E := (\alpha \oplus 0_m, h),$$

where  $h : n + m \rightarrow n + m + q$  is defined as follows:

$$\begin{aligned} h &:= \langle h_1, h_2 \rangle, \quad \text{where} \\ h_1 &= \bar{f} \cdot (\mathbf{1}_n \oplus \beta \oplus 0_q) + (0_n \oplus (a \cdot \beta \cdot g)) \\ h_2 &= 0_n \oplus g. \end{aligned}$$

Note that  $a \cdot \beta \cdot g$  is primitive, by Proposition 9.2. Thus, by Proposition 9.2,  $D \cdot E : k \rightarrow q$  is a primitive normal description, of weight  $n + m$ . Claim:

$$|D \cdot E| = |D| \cdot |E|.$$

*Proof.* First, note that for any morphisms  $F : n \rightarrow n + m + q$  and  $G : m \rightarrow m + q$  in a Conway theory,

$$\langle F, 0_n \oplus G \rangle^\dagger = \langle F^\dagger \cdot \langle G^\dagger, \mathbf{1}_q \rangle, G^\dagger \rangle$$

by the pairing identity,

$$= \langle (F \cdot (\mathbf{1}_n \oplus \langle G^\dagger, \mathbf{1}_q \rangle))^\dagger, G^\dagger \rangle,$$

by the parameter identity. Thus, when  $F = h_1 : n \rightarrow n + m + q$  and  $G = g : m \rightarrow m + q$ , so  $h_2 = 0_n \oplus g$ ,

$$\begin{aligned} (\mathbf{1}_n \oplus 0_m) \cdot h^\dagger &= ((\bar{f} \cdot (\mathbf{1}_n \oplus \beta \oplus 0_q) + 0_n \oplus a \cdot \beta \cdot g) \cdot (\mathbf{1}_n \oplus \langle g^\dagger, \mathbf{1}_q \rangle))^\dagger \\ &= (\bar{f} \cdot (\mathbf{1}_n \oplus \beta \cdot g^\dagger) + (0_n \oplus a \cdot \beta \cdot g^\dagger))^\dagger \\ &= ((\bar{f} + (0_n \oplus a)) \cdot (\mathbf{1}_n \oplus \beta \cdot g^\dagger))^\dagger \\ &= (\bar{f} + 0_n \oplus a)^\dagger \cdot \beta \cdot g^\dagger \\ &= f^\dagger \cdot \beta \cdot g^\dagger. \end{aligned}$$

Thus,

$$\begin{aligned} |D \cdot E| &= (\alpha \oplus 0_m) \cdot h^\dagger \\ &= \alpha \cdot (\mathbf{1}_n \oplus 0_m) \cdot h^\dagger \\ &= \alpha \cdot f^\dagger \cdot \beta \cdot g^\dagger \\ &= |D| \cdot |E|. \end{aligned}$$

**$\mathcal{D}$  is closed under  $\dagger$ .**

Suppose that  $D = (\alpha, f) : k \rightarrow k + p$  has weight  $n$ , so that  $\alpha : k \rightarrow n$  is base and  $f = \bar{f} + (0_n \oplus [a, b]) : n \rightarrow n + k + p$ , where  $a : n \rightarrow k$ ,  $b : n \rightarrow p$  in  $M$ .

Define  $|D^\dagger|$  as  $(\alpha, g)$ , where

$$\begin{aligned} g &= (a \cdot \alpha)^* \cdot (\bar{f} \cdot (\langle \mathbf{1}_n, \alpha \rangle \oplus \mathbf{1}_p) + (0_n \oplus b)) \\ &= ((a \cdot \alpha)^* \cdot \bar{f} \cdot (\langle \mathbf{1}_n, \alpha \rangle \oplus \mathbf{1}_p)) + (0_n \oplus (a \cdot \alpha)^* \cdot b). \end{aligned}$$

We have used the fact that  $M$  has a  $*$  operation, and composition on the left with  $M$ -morphisms distributes over sums.

We claim

$$|D^\dagger| = |D|^\dagger.$$

Indeed,

$$\begin{aligned} |D|^\dagger &= (\alpha \cdot (\bar{f} + (0_n \oplus [a, b])))^\dagger \\ &= \alpha \cdot ((\bar{f} + (0_n \oplus [a, b]))^\dagger \cdot (\alpha \oplus \mathbf{1}_p))^\dagger, \end{aligned}$$

by the composition identity,

$$= \alpha \cdot ((\bar{f} + (0_n \oplus [a, b])) \cdot (\mathbf{1}_n \oplus \alpha \oplus \mathbf{1}_p))^{\dagger\dagger},$$

by the parameter identity,

$$= \alpha \cdot ((\bar{f} + (0_n \oplus [a, b])) \cdot (\mathbf{1}_n \oplus \alpha \oplus \mathbf{1}_p) \cdot (\langle \mathbf{1}_n, \mathbf{1}_n \rangle \oplus \mathbf{1}_p))^\dagger,$$

by the double dagger identity,

$$\begin{aligned} &= (\alpha \cdot (\bar{f} + (0_n \oplus [a, b])) \cdot (\langle \mathbf{1}_n, \alpha \rangle \oplus \mathbf{1}_p))^\dagger \\ &= \alpha \cdot ((\bar{f} \cdot (\langle \mathbf{1}_n, \alpha \rangle \oplus \mathbf{1}_p) + [a \cdot \alpha, b])^\dagger)^\dagger \\ &= \alpha \cdot ((a \cdot \alpha)^* \cdot (\bar{f} \cdot (\langle \mathbf{1}_n, \alpha \rangle \oplus \mathbf{1}_p) + (0_n \oplus b))^\dagger)^\dagger \\ &= \alpha \cdot g^\dagger \\ &= |D|^\dagger. \quad \square \end{aligned}$$

**$\mathcal{D}$  is closed under  $+$ .**

Suppose that  $D = (\alpha, f)$  and  $E = (\beta, g)$  are normal descriptions  $n \rightarrow p$  of weights  $w, s$  respectively. Define the normal description  $D + E = (\gamma, h) : n \rightarrow p$  of weight  $w + s + n$  as follows:

$$\begin{aligned} h &:= \langle h_1, h_2, h_3 \rangle, \quad \text{where} \\ h_1 &:= f \cdot (\mathbf{1}_w \oplus 0_s \oplus 0_n \oplus \mathbf{1}_p) \\ h_2 &:= g \cdot (0_w \oplus \mathbf{1}_s \oplus 0_n \oplus \mathbf{1}_p) \\ h_3 &:= \alpha \cdot h_1 + \beta \cdot h_2 \\ \gamma &:= 0_w \oplus 0_s \oplus \mathbf{1}_n. \end{aligned}$$

Then, by two uses of the pairing identity,

$$\begin{aligned} h^\dagger &= \langle f^\dagger, g^\dagger, \alpha \cdot f \cdot \langle f^\dagger, \mathbf{1}_p \rangle, \beta \cdot g \cdot \langle g^\dagger, \mathbf{1}_p \rangle \rangle \\ &= \langle f^\dagger, g^\dagger, \alpha \cdot f^\dagger + \beta \cdot g^\dagger \rangle, \end{aligned}$$

so that  $|D + E| = \gamma \cdot h^\dagger = |D| + |E|$ . □

**$\mathcal{D}$  is closed under pairing.**

Suppose that  $D = (\alpha, f)$  and  $E = (\beta, g)$  are normal descriptions  $n \rightarrow p$  of weights  $w, s$  respectively. Define the normal description  $\langle D, E \rangle = (\gamma, h) : n \rightarrow p$  of weight  $w + s$  as follows:

$$\begin{aligned} h &:= \langle h_1, h_2 \rangle, \quad \text{where} \\ h_1 &:= f \cdot (\mathbf{1}_w \oplus 0_s \oplus \mathbf{1}_p) \\ h_2 &:= g \cdot (0_w \oplus \mathbf{1}_s \oplus \mathbf{1}_p) \\ \gamma &:= \langle \alpha \cdot (\mathbf{1}_w \oplus 0_s), \beta \cdot (0_w \oplus \mathbf{1}_s) \rangle \end{aligned}$$

Then, by the pairing identity,

$$h^\dagger = \langle f^\dagger, g^\dagger \rangle,$$

so that

$$\begin{aligned} |\langle D, E \rangle| &= \gamma \cdot h^\dagger \\ &= \langle \alpha \cdot f^\dagger, \beta \cdot g^\dagger \rangle \\ &= \langle |D|, |E| \rangle. \quad \square \end{aligned}$$

**$\mathcal{D}$  contains all morphisms in  $M$ .**

For  $[a, b] : n \rightarrow p$  in  $M$ , let  $D$  be the description  $(\alpha, f)$ , where

$$\begin{aligned} \alpha &= \mathbf{1}_n \\ f &= 0_n \oplus [a, b]. \end{aligned}$$

Then, by the left zero identity,

$$\begin{aligned} |D| &= \alpha \cdot (0_n \oplus [a, b])^\dagger \\ &= [a, b]. \end{aligned}$$

It follows that all base morphisms are in  $\mathcal{D}$ .

**$\mathcal{D}$  contains all morphisms in  $\Sigma$ .**

For  $\sigma : 1 \rightarrow k$ , let  $D$  be the description  $(\alpha, f)$ , where

$$\begin{aligned} \alpha &= \mathbf{1}_1 \\ f &= \sigma \cdot (0_1 \oplus \mathbf{1}_k) \\ &= 0_1 \oplus \sigma. \end{aligned}$$

Then, by the left zero identity,  $\sigma = |D|$ . □

We have shown that the behaviors form a subtheory of  $T$  containing  $M \cup \Sigma$ , closed under  $+, \dagger$ . Since Conway grove theories are defined by equations, any subtheory of a Conway grove theory closed under sum and  $\dagger$  will also be a Conway grove theory. This completes the proof.  $\square$

## 10 An application of the Kleene type theorem

In this section we describe an application of the general result given in Theorem 9.4 to formal tree series. But first we give a reformulation of Theorem 9.4 using rational expressions.

**Definition 10.1** *Suppose that  $\Sigma$  is a ranked set and  $S$  is a Conway semiring, or more generally, a semiring equipped with an operation  $*$  :  $S \rightarrow S$ . A rational expression  $1 \rightarrow n$  over  $\Sigma$  with coefficients in  $S$ , or  $(\Sigma, S)$ -rational expression  $1 \rightarrow n$ , for short, is either a symbol  $i_n$ , where  $n \geq 0$  and  $i \in [n]$ , or a symbol in  $\Sigma_n$ , or an expression of one of the following four forms:*

1. Scalar multiplication.  $s \cdot f$ , where  $s \in S$  and  $f$  is a  $(\Sigma, S)$ -rational expression  $1 \rightarrow n$ .
2. Sum.  $f + g$ , where  $f, g$  are  $(\Sigma, S)$ -rational expressions  $1 \rightarrow n$ .
3. Composition.  $f \cdot \langle g_1, \dots, g_m \rangle$ , where  $f$  is a  $(\Sigma, S)$ -rational expression  $1 \rightarrow m$  and  $g_1, \dots, g_m$  are  $(\Sigma, S)$ -rational expressions  $1 \rightarrow n$ . (When  $m = 0$ , we also write  $f \cdot 0_n$ .)
4. Scalar iteration.  $f^\dagger$ , where  $f$  is a  $(\Sigma, S)$ -rational expression  $1 \rightarrow 1 + n$ .

A  $(\Sigma, S)$ -rational expression  $m \rightarrow n$  is an  $m$ -tuple  $\langle f_1, \dots, f_m \rangle$  of  $(\Sigma, S)$ -rational expressions  $f_i : 1 \rightarrow n$ . In particular, for each  $n$ ,  $0_n$  is a  $(\Sigma, S)$ -rational expression  $0 \rightarrow n$ .

**Definition 10.2** *Given a Conway grove theory  $T$  with a sub Conway theory  $M$  which is a matrix theory such that  $M(1,1)$  is (isomorphic to)  $S$ , and given an assignment of a morphism  $\bar{\sigma} : 1 \rightarrow n$  to each letter  $\sigma$  in  $\Sigma_n$ ,  $n \geq 0$ , each  $(\Sigma, S)$ -rational expression  $f : 1 \rightarrow n$  denotes a morphism  $|f| : 1 \rightarrow n$  in  $T$ .*

1. When  $f$  is the expression  $i_n$ , where  $i \in [n]$ ,  $n \geq 0$ , then  $|f|$  is the  $i$ -th distinguished morphism  $i_n : 1 \rightarrow n$  in  $T$ .

2. When  $f = \sigma \in \Sigma_n$ ,  $n \geq 0$ , then  $|f|$  is  $\bar{\sigma}$ .
3. When  $f$  is  $g + h$ , where  $g, h$  are  $(\Sigma, S)$ -rational expressions  $1 \rightarrow n$ , then  $|f|$  is  $|g| + |h|$  in  $T$ .
4. When  $f = g \cdot \langle h_1, \dots, h_m \rangle$ , where  $g$  is a  $(\Sigma, S)$ -rational expression  $1 \rightarrow m$  and each  $h_i$  is a  $(\Sigma, S)$ -rational expression  $1 \rightarrow n$ , then  $|f|$  is the composite  $|g| \cdot \langle |h_1|, \dots, |h_m| \rangle : 1 \rightarrow n$  in  $T$ .
5. When  $f = g^\dagger$ , where  $g$  is a  $(\Sigma, S)$ -rational expression  $1 \rightarrow 1 + n$ , then  $|f|$  is  $|g|^\dagger : 1 \rightarrow n$  in  $T$ .

Suppose that  $f = \langle f_1, \dots, f_n \rangle$  is a  $(\Sigma, S)$ -rational expression  $m \rightarrow n$ . Then we define  $|f|$  as

$$\langle |f_1|, \dots, |f_n| \rangle : m \rightarrow n$$

in  $T$ . We call  $|f|$  the denotation of  $f$  with respect to  $\bar{\Sigma}$ , where  $\bar{\Sigma}_n = \{\bar{\sigma} : \sigma \in \Sigma_n\}$ ,  $n \geq 0$  (or more precisely, with respect to the rank preserving map  $\sigma \mapsto \bar{\sigma}$ ,  $\sigma \in \Sigma$ ).

We may now reformulate Theorem 9.4 as follows:

**Corollary 10.3** *Suppose that  $T$  is a Conway grove theory and equipped with the sub Conway grove theory  $M = \mathbf{Mat}_S$ , where  $S = T(1, 1)$ . Suppose that  $\sigma \mapsto \bar{\sigma}$ ,  $\sigma \in \Sigma$  is a rank preserving map  $\Sigma \rightarrow T$ . Then a morphism  $f : m \rightarrow n$  in  $T$  is the denotation of a  $(\Sigma, S)$ -rational expression  $m \rightarrow n$  with respect to  $\bar{\Sigma}$  iff  $f$  is the behavior  $|D|$  of a normal description  $D : m \rightarrow n$  over  $(S, \bar{\Sigma})$ .*

*Proof.* Immediate from Theorem 9.4 and the fact that in Conway theories, the  $\dagger$ -operation is expressible by the scalar version  $f \mapsto f^\dagger$ ,  $f : 1 \rightarrow 1 + p$  of this operation, and the theory operations. See the pairing identity in Proposition 12.3.  $\square$

**Remark 10.4** *It would be sufficient to allow rational expressions that can be constructed from the symbols  $i_n$  and the letters in  $\Sigma$  by sum, composition, scalar iteration, and left multiplication with the elements of a generating set of  $S$ .*

**Remark 10.5** *It is pointed out in [7] that in theories  $T$  that are Conway grove theories, the  $\dagger$ -operation is equivalent to the following  $\otimes$ -operation. For each  $f : n \rightarrow n + p$ , define*

$$f^\otimes = [(f \cdot (\mathbf{1}_n \oplus \mathbf{0}_n \oplus \mathbf{1}_p)) + (\mathbf{0}_n \oplus \mathbf{1}_n \oplus \mathbf{0}_p)]^\dagger.$$

Thus, when  $T$  is also matrix theory and  $f = a : n \rightarrow n$ , then  $f^\otimes = f^*$ .

By the pairing identity, the  $\dagger$ -operation is expressible by the theory operations and the scalar version of the  $\dagger$ -operation. The same holds true for the  $\otimes$ -operation in Conway grove theories. The scalar  $\otimes$ -operation corresponds to the “ $x_1$ -iteration operation” used in tree automata theory.

Suppose now that  $T$  is the theory  $S\langle\langle \Sigma \rangle\rangle$ , where  $S$  is a Conway semiring, so that  $S\langle\langle \Sigma \rangle\rangle$  is Conway grove theory. Moreover, suppose that  $M$  denotes the sub grove theory of  $S\langle\langle \Sigma \rangle\rangle$  considered in Proposition 8.11, so that  $M$  is isomorphic to the matrix theory  $\mathbf{Mat}_S$ . Since  $S$  is a Conway semiring,  $M$  is a sub Conway grove theory of  $T$ . A normal description  $D = (\alpha, f) : n \rightarrow p$  of weight  $w$  may now be identified with a “weighted  $p$ -ary  $\Sigma$ -tree automaton” over the semiring  $S$  whose set of states is  $[w]$ . For each  $\sigma \in \Sigma_k$  and  $i, i_1, \dots, i_k$  in  $[w]$ , there is a  $\sigma$ -labeled transition from the  $k$ -tuple  $(i_1, \dots, i_k)$  to state  $i$  iff for some nonzero  $s \in S$ , the morphism (i.e., series)  $s \cdot \sigma \cdot \rho$  is a summand of the  $i$ th component  $f_i$  of  $f$ , where  $\rho = \langle i_1, \dots, i_k \rangle$ . In this case,  $s$  is the weight of the transition. (We assume without loss of generality that each  $f_i$  is written as a sum of type 1 and type 2 primitive morphisms  $s \cdot \sigma \cdot \rho$  and  $0_n \oplus (s \cdot \tau)$ , where  $s \neq 0$ , and that any two of these primitive morphisms differ in  $\sigma$ , or in the base morphisms  $\rho$  or  $\tau$ .) The initial and final weight distributions of the automaton are given so that the initial weight of state  $i$  with respect to a variable  $x_j$  with  $j \in [p]$  is the coefficient  $s$  in the summand  $0_n \oplus (s \cdot j_w)$  of the morphism  $f_i$ , if this summand exists, and is 0 otherwise. The final weight of state  $i$  is the  $i$ th component  $\alpha_i$  of the morphism  $\alpha$ . We define the behavior of the  $p$ -ary weighted tree automaton associated with  $D$  as the behavior  $\alpha \cdot f^\dagger$  of the description  $D$ . It is known, cf. [7], that when  $S$  is a continuous semiring as defined by Kuich in [32], or well  $\omega$ -additive as defined in [14], then  $S$  is an iteration semiring and thus a Conway semiring. Moreover, the behavior of a  $p$ -ary weighted  $\Sigma$ -tree automaton as defined in [32] is just the behavior as defined here. Thus, we conclude that Theorem 9.4 implies the Kleene type theorems of [14, 32]. (The rational expressions of [14, 32] are defined in a slightly different but in fact equivalent way.)

While this paper was under review, M. Droste and H. Vogler reported in [21] yet another version of a Kleene-type theorem applicable to formal tree series with coefficients in a commutative and idempotent semiring. Since not every such semiring is a Conway semiring, and since there exist Conway semirings which are neither idempotent nor commutative, the application of Theorem 9.4 described above is independent from the result of [21]. It is usually not possible to define a  $*$ -operation with nice algebraic properties on a semiring, even when the semiring is idempotent and commutative.

## References

- [1] S. Banach, Sur les operations dans les ensembles abstraits et leur applications aux équations intégrales, *Fundamenta Mathematica*, 22(1922), 133–181.
- [2] H. Bekič, Definable operations in general algebras, and the theory of automata and flowcharts, *Technical Report*, IBM Laboratory, Vienna, 1969.
- [3] D. Benson and J. Tiuryn, Fixed points in free process algebras, *Theoretical Computer Science* 63(1989), 275–294.
- [4] J. Berstel, and C. Reutenauer. Recognizable formal power series on trees. *Theoretical Computer Science*, 18(1982), 115–148.
- [5] S.L. Bloom, C.C. Elgot and J.B. Wright, Solutions of the iteration equation and extension of the scalar iteration operation, *SIAM J. Computing*, 9(1980), 26–45.
- [6] S.L. Bloom, C.C. Elgot and J.B. Wright, Vector iteration of pointed iterative theories, *SIAM J. Computing*, 9(1980), 525–540.
- [7] S.L. Bloom and Z. Ésik, *Iteration Theories: The Equational Logic of Iterative Processes*, EATCS Monographs on Theoretical Computer Science, Springer–Verlag, 1993.
- [8] S.L. Bloom and Z. Ésik, Matrix and matricial iteration theories, Part I, *J. of Computer and System Sciences*, 46(1993), 381–408.
- [9] S.L. Bloom and Z. Ésik, Matrix and matricial iteration theories, Part II, *J. of Computer and System Sciences*, 46(1993), 409–439.
- [10] S.L. Bloom and Z. Ésik, The equational logic of fixed points. *Theoretical Computer Science* 179(1997), 1–60.
- [11] S.L. Bloom and Z. Ésik, An extension theorem with an application to formal tree series. *BRICS Report Series*, RS-02-19, May 2002.
- [12] S.L. Bloom, S. Ginali and J. Rutledge, Scalar and vector iteration, *J. Computer and System Sciences*, 14(1977), 251–256.
- [13] S. Bozapalidis. Convex algebras, convex modules and formal power series on trees. *J. Autom. Lang. Comb.* 1 (1996), no. 3, 165–180.
- [14] S. Bozapalidis. Equational elements in additive algebras. *Theory Comput. Systems*, 31 (1999), 1–33.
- [15] L.E.J. Brouwer, Uber Abbildung von Mannigfaltigkeiten, *Math. Annalen*, 71(1910), 97–115.
- [16] J.C. Conway, *Regular Algebra and Finite Machines*, Chapman and Hall, London, 1971.

- [17] J.W. De Bakker and D. Scott, A theory of programs, Technical report, IBM Vienna, 1969.
- [18] J.W. de Bakker and J.I. Zucker, Processes and the denotational semantics of concurrency, *Information and Control*, 54(1982), 70–120.
- [19] J.W. de Bakker and J.N. Kok, Comparative Metric Semantics, *Theoretical Computer Science* 75(1-2), 1990, 14–43.
- [20] J.W. de Bakker and J.H.A. Warmerdam. Four domains for concurrency. *Theoretical Computer Science*, 90(1), 1991, 127–149.
- [21] M. Droste and H. Vogler, A Kleene theorem for weighted tree automata, Technical Report, TUD-FI02-04, TU Dresden, June 2002.
- [22] C.C. Elgot, Monadic computation and iterative algebraic theories, in: *Logic Colloquium 1973, Studies in Logic*, J.C. Shepherdson, editor, volume 80, North Holland, Amsterdam, 1975, 175–230.
- [23] C.C. Elgot, Matricial theories, *J. of Algebra*, 42(1976), 391–421.
- [24] J. Engelfriet and E.M. Schmidt. IO and OI. I. *J. Comput. System Sci.* 15 (1977), 328–353.
- [25] Z. Ésik, Group axioms for iteration. *Information and Computation*, 148(1999), 131–180.
- [26] S.C. Kleene, Representation of events in nerve nets and finite automata. In C.E. Shannon and J. McCarthy, editors, *Automata Studies*, Princeton U. Press, Princeton, NJ, 1956. 3–41.
- [27] P. Freyd, Algebraically complete categories, in: *Category Theory, Como 1990*, LNM vol. 1488, Springer–Verlag, 1991, 95–104.
- [28] P. Freyd, Remarks on algebraically compact categories, in: *Applications of Categories in Computer Science*, London Math. Society Lecture Notes Series, vol. 77, Cambridge University Press, 1992, 95–106.
- [29] D. Krob, Complete systems of B-rational identities, *Theoretical Computer Science*, 89(1991), 207–343.
- [30] A. Bonnier-Rigny and D. Krob. A complete system of identities for one-letter rational expressions with multiplicities in the tropical semiring. *Theoret. Comput. Sci.* 134 (1994), no. 1, 27–50.
- [31] W. Kuich. Automata and languages generalized to  $\omega$ -continuous semirings, *Theoret. Comp. Sci.*, 79(1991), 137–150.

- [32] W. Kuich. Formal power series over trees. In *Proc. 3rd International Conf. Developments in Language Theory* (S. Bozapalidis, ed), Aristotle University of Thessaloniki, 1998, 61–101.
- [33] F.W. Lawvere. Functorial semantics of algebraic theories. In: *Proceedings of the National Academy of Sciences USA*, 50(1963), 869–873.
- [34] D.J. Lehmann and M.B. Smyth. Algebraic specification of data types: a synthetic approach. *Math. Systems Theory*, 14 (1981), no. 2, 97–139.
- [35] R. Milner. A complete inference system for a class of regular behaviors. *J. Comp. System Sci.*, 28 (1984), 439–466.
- [36] D. Niwinski, On fixed-point clones, in: *Proc. ICALP '86*, LNCS 226, Springer–Verlag, 1986, 464–473.
- [37] G.D. Plotkin, *Domains*, Lecture Notes, Department of Computer Science, University of Edinburgh, 1983.
- [38] A. Salomaa, Two complete axioms systems for the algebra of regular events, *J. of the ACM*, 13(1966), 158–169.
- [39] A. Salomaa, On regular expressions and regular canonical systems, *Math. Syst. Theory*, 2(1968), 341–355.
- [40] D.S. Scott. Data types as lattices. *SIAM J. Computing*, 5(3), 522–587 (1976).
- [41] A.K. Simpson, A characterisation of the least fixed point operator by dinaturality, *Theoretical Computer Science*, 118(1993), 301–314.
- [42] M.B. Smyth and G.D. Plotkin, The category theoretic solution of recursive domain equations, *SIAM Journal of Computing*, 11(1982), 761–783.
- [43] A. Simpson and G. Plotkin. Complete Axioms for Categorical Fixed-point operations. In: *Proceedings of the 15th Annual IEEE Logic in Computer Science Colloq.*, 30-41, (2000).
- [44] A. Tarski, A lattice-theoretical fixed point theorem and its applications. *Pacific J. Math.*, 5(1955), 285–309.

## 11 Theories and trees

This section is intended as a quick review of the basic notions. We have lifted a great deal from our tutorial paper [10].

A **Lawvere algebraic theory**  $T$ , or just **theory** for short, is a category whose objects are the nonnegative integers in which the object  $n$  is the copower of the object 1 with itself  $n$ -times. (Sometimes Lawvere theories are defined dually with  $n$  being the  $n$ th power of the object 1.) See [33].

We may elaborate the definition somewhat by specifying the copower injections  $i_n : 1 \rightarrow n$ , for  $i \in [n]$ . Thus, the official definition of theory is the following.

**Definition 11.1** *A theory  $T$  is a category whose objects are the nonnegative integers, which has, for each  $n \geq 0$ ,  $n$  “distinguished morphisms”  $i_n : 1 \rightarrow n$ ,  $i \in [n]$ , with the property that for any  $p \geq 0$  and any family of morphisms  $g_i : 1 \rightarrow p$ ,  $i \in [n]$ , there is a unique morphism  $g : n \rightarrow p$  such that*

$$i_n \cdot g = g_i, \quad i \in [n].$$

The uniquely determined morphism  $g$  is denoted  $\langle g_1, \dots, g_n \rangle$ , and is called the “tupling” of the  $g_i$ . Note that tupling determines a bijection

$$\overbrace{T(1, p) \times \dots \times T(1, p)}^n \rightarrow T(n, p),$$

so that one may specify a morphism  $f : n \rightarrow p$  by giving its components  $i_n \cdot f : 1 \rightarrow p$ ,  $i \in [n]$ . We assume that when  $i = n = 1$ , the distinguished morphism  $i_n$  is the identity morphism  $1 \rightarrow 1$ , so that  $\langle g \rangle = g$ , for any  $g : 1 \rightarrow p$ . In any theory we write the identity morphism  $p \rightarrow p$  as  $\mathbf{1}_p : p \rightarrow p$ . Note that  $\mathbf{1}_p = \langle 1_p, \dots, 1_p \rangle$ , the tupling of the  $p$  distinguished morphisms  $1 \rightarrow p$ .

The coproduct property implies that the object 0 is an initial object: there is a unique morphism  $0 \rightarrow n$ , for each  $n \geq 0$ . We denote this morphism by

$$0_n : 0 \longrightarrow n.$$

In any theory, a morphism with source 1 is called a **scalar morphism**.

If  $T$  and  $T'$  are theories, a **theory morphism**  $T \rightarrow T'$  is a functor which preserves the objects and distinguished morphisms. It follows that any theory morphism preserves tupling. (Morphisms  $n \rightarrow p$  in some theory  $T$  might also legitimately be called “theory morphisms”, but we call them  $T$ -morphisms, or just “morphisms”.)

There is an “initial theory” **Tot**, the theory of all total functions  $[n] \rightarrow [p]$ ,  $n, p \geq 0$ . If  $T$  is any theory, there is a unique theory morphism **Tot**  $\rightarrow T$  under which the function  $f : [n] \rightarrow [p]$  is mapped to the  $T$ -morphism

$$\bar{f} := \langle f(1)_p, \dots, f(n)_p \rangle : n \longrightarrow p.$$

The morphisms  $\bar{f}$  determined by functions are called **base morphisms**.

A theory is *trivial* if each hom-set  $T(n, p)$  has at most one morphism. If  $T$  is not trivial, the unique theory morphism from **Tot** is injective and we usually assume without comment that **Tot** is a subtheory of every theory.

It is useful to introduce the names

$$\begin{aligned}\kappa : n &\longrightarrow n + p \\ \lambda : p &\longrightarrow n + p\end{aligned}$$

for the base morphisms which correspond to the inclusion function  $[n] \rightarrow [n + p]$  and translated inclusion function  $[p] \rightarrow [n + p]$ . Using these base morphisms, we introduce two derived operations on theories: *pairing*, which is a function

$$\begin{aligned}T(n, p) \times T(m, p) &\longrightarrow T(n + m, p) \\ f, g &\mapsto \langle f, g \rangle\end{aligned}$$

for all  $n, m, p \geq 0$ , and *separated sum*, which is an operation

$$\begin{aligned}T(n, p) \times T(m, q) &\longrightarrow T(n + m, p + q) \\ f, g &\mapsto f \oplus g.\end{aligned}$$

The definition of pairing is the following. For  $f : n \rightarrow p$ ,  $g : m \rightarrow p$  in  $T$ ,  $i \in [n + m]$ ,

$$i_{n+m} \cdot \langle f, g \rangle := \begin{cases} i_n \cdot f & \text{if } i \in [n] \\ j_m \cdot g & \text{if } i = n + j, j \in [m]. \end{cases}$$

The definition of separated sum makes use of  $\kappa : p \rightarrow p + q$ ,  $\lambda : q \rightarrow p + q$ , when  $f : n \rightarrow p$ ,  $g : m \rightarrow q$ .

$$i_{n+m} \cdot (f \oplus g) := \begin{cases} i_n \cdot f \cdot \kappa & \text{if } i \in [n] \\ j_m \cdot g \cdot \lambda & \text{if } i = n + j, j \in [m]. \end{cases}$$

There is an interesting calculus of theory terms built from the operations of pairing and separated sum. For example, the following equations are valid in all theories,

when sources and targets are appropriate:

$$\begin{aligned}
\kappa \cdot \langle f, g \rangle &= f \\
\lambda \cdot \langle f, g \rangle &= g \\
\langle \kappa \cdot f, \lambda \cdot f \rangle &= f \\
\langle f, \langle g, h \rangle \rangle &= \langle \langle f, g \rangle, h \rangle \\
\langle f, g \rangle \cdot h &= \langle f \cdot h, g \cdot h \rangle \\
f \oplus (g \oplus h) &= (f \oplus g) \oplus h \\
(f \oplus g) \cdot \langle h, k \rangle &= \langle f \cdot h, g \cdot k \rangle \\
(f \oplus g) \cdot (h \oplus k) &= (f \cdot h) \oplus (g \cdot k)
\end{aligned} \tag{13}$$

We can express both  $\kappa : p \rightarrow p + q$  and  $\lambda : q \rightarrow p + q$  using the separated sum operation:

$$\begin{aligned}
\kappa &= \mathbf{1}_p \oplus 0_q \\
\lambda &= 0_p \oplus \mathbf{1}_q
\end{aligned}$$

We note some further rules for the zero-morphisms  $0_p$ , assuming the expressions are meaningful.

$$\begin{aligned}
\langle 0_p, f \rangle &= f = \langle f, 0_p \rangle \\
\langle f, g \rangle \oplus 0_p &= \langle f \oplus 0_p, g \oplus 0_p \rangle \\
0_p \oplus \langle f, g \rangle &= \langle 0_p \oplus f, 0_p \oplus g \rangle \\
f \oplus 0_q &= f \cdot (\mathbf{1}_p \oplus 0_q) \\
0_q \oplus f &= f \cdot (0_q \oplus \mathbf{1}_p)
\end{aligned}$$

*Node Labeled Trees.* Suppose that  $\Sigma = \Sigma_n, n \geq 0$ , is a ranked set, and  $X = \{x_1, \dots, x_n, \dots\}$  is a countably infinite set of “variables” disjoint from each set  $\Sigma_n$ . We write  $X_p$  for the set  $\{x_1, \dots, x_p\}$ .

A (possibly infinite)  $\Sigma$ -**tree**  $t : 1 \rightarrow p$  is a rooted, finitely branching tree whose leaves are labeled by letters in  $X_p \cup \Sigma_0$ , and whose interior nodes of outdegree  $n \geq 1$  are labeled by letters in  $\Sigma_n$ . Any such tree may be modeled as a partial function  $t : [\omega]^* \rightarrow \cup_{n \geq 0} \Sigma_n \cup X_p$ , where  $[\omega]$  denotes the positive integers, subject to the following conditions:

- The domain of  $t$  is a nonempty, prefix closed subset of  $[\omega]^*$ .
- If  $t(u) \in \Sigma_n$ , for some  $u \in [\omega]^*$  and  $n \geq 0$ , then for any  $i \in [\omega]$ ,  $t(ui)$  is defined iff  $i \in [n]$ .

- If  $t(u) \in X_p$ , then  $t(ui)$  is not defined for any  $i \in [\omega]$ .

Let  $\Sigma \text{TR}(1, p)$  denote the set of all such trees. The subset of trees in  $\Sigma \text{TR}(1, p)$  with a finite domain is denoted  $T_\Sigma(X_p)$ . The tree  $1 \rightarrow p$  whose root is a leaf labeled  $x_i$ ,  $i \in [p]$  is denoted  $i_p$ .

One may compose a tree  $t$  in  $\Sigma \text{TR}(1, p)$  with a  $p$ -tuple of trees  $(g_1, \dots, g_p)$  in  $\Sigma \text{TR}(1, n)$  obtaining the tree denoted

$$t \cdot \langle g_1, \dots, g_p \rangle$$

in  $\Sigma \text{TR}(1, n)$  by attaching a copy of the tree  $g_i$  to each leaf of  $t$  labeled  $x_i$ ,  $i \in [p]$ . Note that

$$\begin{aligned} t \cdot \langle 1_p, \dots, p_p \rangle &= t \\ i_p \cdot \langle g_1, \dots, g_p \rangle &= g_i, \quad i \in [p]. \end{aligned}$$

We let  $\Sigma \text{TR}(n, p)$  denote the collection of  $n$ -tuples of trees in  $\Sigma \text{TR}(1, p)$ .

The  $\Sigma$ -trees form a theory  $\Sigma \text{TR}$ , with  $n$ -tuples of  $\Sigma$ -trees in  $\Sigma \text{TR}(1, p)$  as morphisms  $n \rightarrow p$ . A letter  $\sigma \in \Sigma_n$  is identified with the “atomic tree”

$$\eta(\sigma) : 1 \longrightarrow n$$

whose root is labeled  $\sigma$ ; the  $n$ -successors of the root are leaves, labeled in order  $x_1, \dots, x_n$ .

For  $i \in [p]$ , the letter  $x_i$  is identified with the tree  $\eta(x_i)$ , consisting only of a root labeled  $x_i$ .

The theory operation of tupling  $\langle g_1, \dots, g_n \rangle$  when applied to the  $n$ -tuple  $(g_1, \dots, g_n)$  in  $\Sigma \text{TR}(1, p)$  produces the  $n$ -tuple  $(g_1, \dots, g_n)$ , now considered a morphism  $g : n \rightarrow p$ . The composite of  $\langle t_1, \dots, t_k \rangle : k \rightarrow n$  with this  $g$  is forced by (13) above to be defined as

$$\langle t_1, \dots, t_k \rangle \cdot g = \langle t_1 \cdot g, \dots, t_k \cdot g \rangle, \quad t_i : 1 \longrightarrow n, \quad g : n \longrightarrow p.$$

Note that if  $t$  and the  $g_i$  have finite domains, so does  $t \cdot \langle g_1, \dots, g_p \rangle$ . Thus, the finite trees form a subtheory of  $\Sigma \text{TR}$ , denoted  $T_\Sigma$ .

If  $t : [\omega^*] \rightarrow (\bigcup_n \Sigma_n) \cup X_p$  is a tree and  $u \in [\omega^*]$ , we write  $t_u$  for the tree

$$t_u(v) := t(uv), \quad v \in [\omega^*].$$

A tree  $t$  is regular if the set  $\{t_u : u \in [\omega^*]\}$  is finite. The collection of regular trees determines another subtheory of  $\Sigma \text{TR}$ , denote  $\Sigma \text{tr}$ .

Recall that when  $\Sigma$  is ranked set, a  $\Sigma$ -**algebra**  $B$  is a set equipped with a function

$$\sigma_B : B^n \longrightarrow B$$

for each  $\sigma \in \Sigma_n$ . If  $B, C$  are  $\Sigma$ -algebras, a  $\Sigma$ -**algebra morphism**  $h : B \rightarrow C$  is a function such that for all  $n \geq 0$  and all  $\sigma \in \Sigma_n$ ,

$$h(\sigma_B(b_1, \dots, b_n)) = \sigma_C(h(b_1), \dots, h(b_n)),$$

$b_i \in B$ .

For each set  $p \geq 0$ , the sets  $\Sigma \text{TR}(1, p)$  and  $T_\Sigma(1, p)$  are  $\Sigma$ -algebras, where for  $\sigma \in \Sigma_n$ ,  $t_i \in \Sigma \text{TR}(1, p)$ ,

$$\sigma(t_1, \dots, t_n) := \sigma \cdot \langle t_1, \dots, t_n \rangle.$$

We take it that the following facts are well-known.

**Theorem 11.2 (Free theories)** *The theory  $T_\Sigma$  is the algebraic theory freely generated by  $\Sigma$ , i.e., for any algebraic theory  $T$  and any function  $h$  mapping  $\Sigma_n \rightarrow B(1, n)$ , for each  $n \geq 0$ , there is a unique theory morphism  $h^\# : T_\Sigma \rightarrow T$  such that*

$$h^\#(\eta(\sigma)) = h(\sigma),$$

for all  $\sigma \in \Sigma_n$ ,  $n \geq 0$ .

## 12 Conway and iteration theories

**Definition 12.1** *A preiteration theory is a theory equipped with a dagger operation*

$$\begin{array}{ccc} T(n, n+p) & \longrightarrow & T(n, p) \\ f & \mapsto & f^\dagger \end{array}$$

for each  $n, p \geq 0$ . This operation need not satisfy any particular properties. A **morphism**  $\varphi : T \rightarrow T'$  **of preiteration theories** is a theory morphism which preserves the dagger operation, i.e.,

$$\varphi(f^\dagger) = n\varphi(f)^\dagger, \quad f : n \longrightarrow n+p.$$

**Notation.** In any preiteration theory, we let  $\perp_{n,p} : n \rightarrow p$  abbreviate  $(\mathbf{1}_n \oplus 0_p)^\dagger$ , and when  $p = 0$ , we write just  $\perp_n$ , and when  $n = 1$ , just  $\perp$ .

**Definition 12.2** *A Conway theory is a preiteration theory satisfying the following equational axioms:*

1. PARAMETER IDENTITY

$$(f \cdot (\mathbf{1}_n \oplus g))^\dagger = f^\dagger \cdot g,$$

*all  $f : n \rightarrow n + p$ ,  $g : p \rightarrow q$ . When  $n = 1$ , this equation is called the scalar parameter identity.*

2. COMPOSITION IDENTITY

$$(f \cdot \langle g, 0_n \oplus \mathbf{1}_p \rangle)^\dagger = f \cdot \langle (g \cdot \langle f, 0_m \oplus \mathbf{1}_p \rangle)^\dagger, \mathbf{1}_p \rangle,$$

*all  $f : n \rightarrow m + p$ ,  $g : m \rightarrow n + p$ . When  $m = n = 1$ , this equation is the scalar composition identity.*

3. DOUBLE DAGGER IDENTITY

$$f^{\dagger\dagger} = (f \cdot (\langle \mathbf{1}_n, \mathbf{1}_n \rangle \oplus \mathbf{1}_p))^\dagger,$$

*all  $f : n \rightarrow n + n + p$ . When  $n = 1$ , this equation is called the scalar double dagger identity.*

**A morphism of Conway theories** is a preiteration theory morphism.

Since we have defined Conway theories by equations, they form a variety of preiteration theories. We give several more non-trivial equations that hold in Conway theories.

**Proposition 12.3** *Each Conway theory  $T$  satisfies the following identities.*

1. (ELGOT) FIXED POINT IDENTITY

$$f^\dagger = f \cdot \langle f^\dagger, \mathbf{1}_p \rangle,$$

*all  $f : n \rightarrow n + p$ . When  $n = 1$ , this identity is called the scalar fixed point identity.*

## 2. LEFT ZERO IDENTITY

$$(0_n \oplus f)^\dagger = f,$$

all  $f : n \rightarrow p$ . When  $n = 1$ , this identity is called the scalar left zero identity.

## 3. RIGHT ZERO IDENTITY

$$(f \oplus 0_q)^\dagger = f^\dagger \oplus 0_q,$$

all  $f : n \rightarrow n + p$ . When  $n = 1$ , this identity is called the scalar right zero identity.

## 4. PAIRING IDENTITY

$$\langle f, g \rangle^\dagger = \langle f^\dagger \cdot \langle h^\dagger, \mathbf{1}_p \rangle, h^\dagger \rangle,$$

all  $f : n \rightarrow n + m + p$ ,  $g : m \rightarrow n + m + p$ , where

$$h = g \cdot \langle f^\dagger, \mathbf{1}_{m+p} \rangle : m \longrightarrow m + p.$$

## 5. PERMUTATION IDENTITY

$$(\pi \cdot f \cdot (\pi^{-1} \oplus \mathbf{1}_p))^\dagger = \pi \cdot f^\dagger,$$

for all  $f : n \rightarrow n + p$  and for all base permutations  $\pi : n \rightarrow n$ . Here  $\pi^{-1}$  denotes the inverse of  $\pi$ .

(There are several other axiomatizations of Conway theories, see [7].)

## 13 Iteration theories

Suppose that  $G = (G, \circ)$  is a finite group on the set  $[n]$ . In each theory  $T$ , we associate with  $G$  the base morphisms  $\rho_i^G : n \rightarrow n$ ,  $i \in [n]$ , defined by:

$$\begin{aligned} j_n \cdot \rho_i^G &:= (i \circ j)_n, \quad \text{all } j \in [n], \text{ i.e.,} \\ \rho_i^G &:= \langle (i \circ 1)_n, (i \circ 2)_n, \dots, (i \circ n)_n \rangle. \end{aligned}$$

From now on, we write just  $ij$  instead of  $i \circ j$ . The morphism  $(ij)_n$  is the corresponding base morphism  $1 \rightarrow n$ . The morphisms  $\rho_i^G$ , denoted sometimes just  $\rho_i$ , are called the **base morphisms associated with the group  $G$** .

We define a morphism  $g_G : n \rightarrow n + p$  for each scalar  $g : 1 \rightarrow n + p$ .

$$g_G := \langle g \cdot (\rho_1^G \oplus \mathbf{1}_p), \dots, g \cdot (\rho_n^G \oplus \mathbf{1}_p) \rangle : n \longrightarrow n + p,$$

where  $\tau_n : n \rightarrow 1$  is the unique base morphism.

**Definition 13.1** *The group-identity associated with  $G$  is the equation*

$$g_G^\dagger = \tau_n \cdot (g \cdot (\tau_n \oplus \mathbf{1}_p))^\dagger, \quad g : 1 \longrightarrow n + p. \quad (14)$$

**Example 13.2** *Suppose that  $G$  is the group of order 3. Then the identity associated with  $G$  is the equation*

$$\langle f, f \cdot (\langle 2_3, 3_3, 1_3 \rangle \oplus \mathbf{1}_p), f \cdot (\langle 3_3, 1_3, 2_3 \rangle \oplus \mathbf{1}_p) \rangle^\dagger = \tau_3 \cdot (f \cdot (\tau_3 \oplus \mathbf{1}_p))^\dagger,$$

where  $f : 1 \rightarrow 3 + p$ .

**Definition 13.3** *An iteration theory is a Conway theory which satisfies all identities associated with finite (simple) groups. A morphism of iteration theories is a preiteration theory morphism.*

Free iteration theories are also tree theories.

Let  $\Sigma$  be a signature, and let  $\perp$  be a letter not in any set  $\Sigma_n$ .  $\Sigma_\perp$  is the signature obtained from  $\Sigma$  by adding the letter  $\perp$  to  $\Sigma_0$ . In the tree theory  $\Sigma_\perp \text{ tr}$ ,  $\mathbf{1}_1^\dagger = \perp$ .

**Theorem 13.4** [7] *For any signature  $\Sigma$ , the theory  $\Sigma_\perp \text{ tr}$  of regular  $\Sigma_\perp$ -trees is the iteration theory freely generated by  $\Sigma$ ; for any iteration theory  $T$  and any function  $h : \Sigma \rightarrow T$  which maps  $\sigma \in \Sigma_n$  to a morphism  $h(\sigma) : 1 \rightarrow n$  in  $T$ , all  $n \geq 0$ , there is a unique iteration theory morphism  $h^\# : \Sigma_\perp \text{ tr} \rightarrow T$  such that*

$$h(\sigma) = h^\#(\eta(\sigma)),$$

for all  $\sigma \in \Sigma_n$ ,  $n \geq 0$ .

**Example 13.5** *Let  $L = (L, \leq)$  be a complete (or just  $\omega$ -complete) poset. Let  $T_L$  denote the theory whose morphisms  $1 \rightarrow n$  are the continuous (i.e., sup preserving) functions  $L^n \rightarrow L$ ; the composite of  $f : 1 \rightarrow n$  with  $\langle g_1, \dots, g_n \rangle : n \rightarrow p$  is the function*

$$L^p \xrightarrow{\langle g_1, \dots, g_n \rangle} L^n \xrightarrow{f} L.$$

For each continuous  $f : L^{n+p} \rightarrow L^n$  and each  $a \in L^p$  there is a least  $b \in L^n$  such that  $b = f(b, a)$ . This  $b$  is denoted  $\mu b. b = f(b, a)$ . It was shown in several places that  $T_L$  is an iteration theory, where, for  $f : n \rightarrow n + p$ , (i.e.,  $f : L^{n+p} \rightarrow L^n$ )

$$f^\dagger(a) = \mu b. b = f(b, a).$$

There is a number of classes of theories whose identities are captured by the axioms of iteration theories.

**Theorem 13.6** [7] *An identity holds in all iteration theories iff it holds in all theories  $T_L$ , for  $\omega$ -complete posets  $L$ .*

Other representative classes are theories in which the fixed point operation is a “unique fixed point”, or alternatively, an “initial fixed point”. See [10] for a thorough account.

A sufficient condition that a Conway theory satisfies the group identities is that it satisfies the implication we have called the **weak functorial implication**. (Actually, there is one such implication for each  $n \geq 1$ .) A preiteration theory satisfies the weak functorial implication if, whenever the square

$$\begin{array}{ccc}
 n & \xrightarrow{f} & n + p \\
 \tau_n \downarrow & & \downarrow \tau_n \oplus \mathbf{1}_p \\
 1 & \xrightarrow{g} & 1 + p
 \end{array}$$

commutes, then so does

$$\begin{array}{ccc}
 n & \xrightarrow{f^\dagger} & p \\
 \tau_n \downarrow & \nearrow g & \\
 1 & & 
 \end{array}$$

Here,  $\tau_n : n \rightarrow 1$  is the unique base morphism.

For example, when  $L$  is an  $\omega$ -complete poset,  $T_L$  satisfies the weak functorial implication, since

$$f^\dagger = \sup_k f^{(k)},$$

where the powers of  $f$  are defined by:

$$\begin{aligned}
 f^{(1)} &= f \cdot (\perp_n \oplus \mathbf{1}_p) \\
 f^{(k+1)} &= f \cdot \langle f^{(k)}, \mathbf{1}_p \rangle.
 \end{aligned}$$

Similarly,  $g^\dagger = \sup g^{(k)}$ . But, for each  $k > 0$ ,

$$f^{(k)} = \tau_n \cdot g^{(k)}.$$